



Complete Summary

GUIDELINE TITLE

Low back disorders.

BIBLIOGRAPHIC SOURCE(S)

Hegmann KT, ed. Low back disorders. In: Glass LS, editor(s). Occupational medicine practice guidelines: evaluation and management of common health problems and functional recovery in workers. 2nd ed. Elk Grove Village (IL): American College of Occupational and Environmental Medicine (ACOEM); 2007. p. 366. [1310 references]

GUIDELINE STATUS

This is the current release of the guideline.

This guideline updates a previous version: Low back complaints. In: Glass LS, editor(s). Occupational medicine practice guidelines: evaluation and management of common health problems and functional recovery in workers. 2nd ed. Elk Grove Village (IL): American College of Occupational and Environmental Medicine (ACOEM); 2004. p. 286-326.

The ACOEM *Guidelines* are currently being updated on a 3-year rolling process.

** REGULATORY ALERT **

FDA WARNING/REGULATORY ALERT

Note from the National Guideline Clearinghouse: This guideline references a drug(s) for which important revised regulatory information has been released.

- [December 12, 2007, Carbamazepine](#): The U.S. Food and Drug Administration (FDA) has provided recommendations for screening that should be performed on specific patient populations before starting treatment with carbamazepine.

COMPLETE SUMMARY CONTENT

** REGULATORY ALERT **

SCOPE

METHODOLOGY - including Rating Scheme and Cost Analysis

RECOMMENDATIONS

EVIDENCE SUPPORTING THE RECOMMENDATIONS

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

CONTRAINDICATIONS

QUALIFYING STATEMENTS

SCOPE

DISEASE/CONDITION(S)

Low back disorders

GUIDELINE CATEGORY

Diagnosis
Evaluation
Management
Treatment

CLINICAL SPECIALTY

Family Practice
Internal Medicine
Orthopedic Surgery
Physical Medicine and Rehabilitation
Preventive Medicine
Surgery

INTENDED USERS

Advanced Practice Nurses
Physician Assistants
Physicians
Utilization Management

GUIDELINE OBJECTIVE(S)

- To update the 2004 American College of Occupational and Environmental Medicine's (ACOEM's) Guidelines on Low Back Complaints
- To help improve or restore the health of those workers who incur occupationally related illnesses or injuries
- To present essential evidence-based information to address the injured worker's functional impairment and safely return him or her to work

TARGET POPULATION

Adults with potentially work-related low back disorders seen in primary care settings

INTERVENTIONS AND PRACTICES CONSIDERED

Note from the National Guideline Clearinghouse (NGC): The following general clinical measures were considered. Refer to the "Major Recommendations" section of this summary and the original guideline document for information regarding which specific interventions and practices under these general headings are recommended, optional, or not recommended by the American College of Occupational and Environmental Medicine.

1. Diagnostic testing
2. Medications
3. Orthotics and immobilization
4. Physical treatment methods
5. Activity modification and exercise
6. Injections
7. Surgical considerations
8. Rehabilitation/behavioral/education

MAJOR OUTCOMES CONSIDERED

Missed work days

METHODOLOGY

METHODS USED TO COLLECT/SELECT EVIDENCE

Hand-searches of Published Literature (Secondary Sources)
Searches of Electronic Databases

DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE

Note from the National Guideline Clearinghouse (NGC): The American College of Occupational and Environmental Medicine contracted the Work Loss Data Institute to provide medical library research services.

Disability-Duration Data

This edition includes disability-duration data that have been extracted from National Health Interview Survey data. Only data from interviews with individuals without workers' compensation claims has been included.

2007 Update

Under the ACOEM new methodology, the process begins with the systematic identification of high-quality original research studies on a topic. References are identified from a number of national and international databases of original research. Additional references are identified through an exhaustive "hand search" of the literature by trained health science researchers.

NUMBER OF SOURCE DOCUMENTS

Not stated

METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE

Expert Consensus
Weighting According to a Rating Scheme (Scheme Given)

RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE

Strength of Evidence Ratings

A: Strong evidence-base: Two or more high-quality studies.¹

B: Moderate evidence-base: At least one high-quality study or multiple moderate-quality studies² relevant to the topic and the working population.

C: Limited evidence-base: At least one study of moderate quality.

I: Insufficient evidence: Evidence is insufficient or irreconcilable.

¹For therapy and prevention, randomized controlled trials (RCTs) with narrow confidence intervals and minimal heterogeneity.

For diagnosis and screening, cross sectional studies using independent gold standards.

For prognosis, etiology or harms, prospective cohort studies with minimal heterogeneity.

²For therapy and prevention, a well-conducted review of cohort studies.

For prognosis, etiology or harms, a well-conducted review of retrospective cohort studies or untreated control arms of RCTs.

METHODS USED TO ANALYZE THE EVIDENCE

Review of Published Meta-Analyses
Systematic Review with Evidence Tables

DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE

Contributors reviewed at least one chapter each and reviewed the relevant medical literature that had been published since the creation of the original Guidelines in 1997.

2007 Update

As part of the update process, American College of Occupational and Environmental Medicine (ACOEM) adopted a new more meticulous strength-of-evidence rating methodology. The enhanced methodology incorporates the highest scientific standards for reviewing evidence-based literature, thus ensuring the most rigorous, reproducible, and transparent occupational health guidelines available.

Studies are graded for actual design and for execution of that design and the subsequent analyses of results. Evidence with the highest available ranking—e.g., all randomized controlled trials (RCTs) or randomized crossover trials for

treatment studies—is selected. Each article that meets inclusion criteria is reviewed and critically appraised.

As an example, RCTs that meet inclusion criteria are scored on 11 criteria. Each criterion is scored 0.0, 0.5, or 1.0. These individual ratings are summed up, resulting in an overall rating that ranges from 0 to 11.

The rating for each article is then converted into a quality grade—low quality (0-3.5), moderate quality (4.0-7.5), or high quality (8.0-11.0).

| Criteria | Rating Description |
|--------------------------------|--|
| Randomization | Assessment of the degree that randomization was both reported to have been performed and successfully achieved through analyses of comparisons of variables between the two groups. |
| Treatment Allocation Concealed | Concealment of the allocation scheme from all involved, not just the patient. |
| Baseline Comparability | Measurement of how well the baseline groups are comparable (e.g., age, gender, disease duration, prior treatment). |
| Patient Blinded | Blinding of the patient/subject to the treatment administered. |
| Provider Blinded | Blinding of the provider to the treatment administered. |
| Assessor Blinded | Blinding of the assessor to the treatment administered. |
| Controlled for Co-intervention | The degree to which the study design controlled for multiple interventions (e.g., a combination of stretching exercises and anti-inflammatory medication or mention of not using other treatments during the study). |
| Compliance Acceptable | Measurement of the degree of non-compliance. |
| Dropout Rate | Measurement of the drop-out rate. |
| Timing of Assessments | Assessment of whether the timing of measurements of effects is the same between treatment groups. |
| Analyzed by Intention to Treat | Ascertainment of whether the study was analyzed with an intent-to-treat analysis. |

While literature searches also seek systematic reviews and meta-analyses, on critical appraisal very few of these secondary studies are truly systematic as the term is used in the evidence-based medicine literature. Most typically, there are errors in analyses or interpretation. For this reason, ACOEM relies primarily on the original literature as the source for its evidence syntheses and recommendations.

Acceptable studies are abstracted into evidence tables that include details of study methods, outcomes, and statistical analyses. Research staff then use the tables to grade the strength of evidence in order to draft specific clinical practice recommendations that will be combined into collective evidence-based guidelines. Evidence is drawn almost entirely from original research studies.

METHODS USED TO FORMULATE THE RECOMMENDATIONS

Expert Consensus

DESCRIPTION OF METHODS USED TO FORMULATE THE RECOMMENDATIONS

In reviewing or revising recommendations, the expert Panels review the articles, evidence tables, and strength-of-evidence ratings (A, B, C, or I). Panels discuss recommendations for diagnosis or treatment based on the critically appraised body of evidence using a "best evidence" approach. In addition to critically appraised evidence, "first principles" of medical logic and ethics are observed in formulating recommendations:

- Imaging or testing should generally be done to confirm a clinical impression.
- Tests should affect the course of treatment.
- Treatments should improve on the natural history of the disorder, which in many cases is recovery without treatment.
- Invasive treatment should be preceded by adequate conservative treatment and may be performed if conservative treatment does not improve the health problem.
- The more invasive and permanent, the more caution should be exerted in considering invasive tests or treatments and the stronger should be the evidence of efficacy.
- The more costly the test or intervention, the more caution should be generally exerted prior to ordering the test or treatment and the stronger should be the evidence of efficacy.
- Testing/treatment decisions should be a collaboration between the clinician and patient with full disclosure of benefits and risks.
- Treatment should not create dependence or functional disability.

Health benefits, side effects, and risks are explicitly considered and discussed in formulating recommendations. Benefits should significantly exceed risks. Each recommendation specifies the clinical problem to which it relates and is linked to the relevant higher quality available evidence. Consensus recommendations, following the first principles above, are formulated when there is either a lack of quality evidence or the available evidence substantially conflicts.

RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS

The American College of Occupational and Environmental Medicine (ACOEM) evidence-based recommendations are explicitly classified as follows:

| Recommendation Category | Evidence Rating | Description of Category |
|--------------------------------|------------------------|---|
| Strongly Recommended | A | The intervention is strongly recommended for appropriate* patients. The intervention improves important health and functional outcomes based on high quality evidence, and the Evidence-based Practice Panel (EBPP) |

| Recommendation Category | Evidence Rating | Description of Category |
|---|-----------------|--|
| | | concludes that benefits substantially outweigh harms and costs. |
| Moderately Recommended | B | The intervention is recommended for appropriate patients. The intervention improves important health and functional outcomes based on moderate quality evidence that benefits substantially outweigh harms and costs. |
| Recommended | C | The intervention is recommended for appropriate patients. There is limited evidence that the intervention may improve important health and functional benefits. |
| Insufficient - Recommended (Consensus-based) | I | The intervention is recommended for appropriate patients and has nominal costs and essentially no potential for harm.** The EBPP feels that the intervention constitutes best medical practice to acquire or provide information in order to best diagnose and treat a health condition and restore function in an expeditious manner. The EBPP believes based on the body of evidence, first principles, and/or collective experience that patients are best served by these practices, although the evidence is insufficient for an evidence-based recommendation. |
| Insufficient - No Recommendation (Consensus-based) | I | The evidence is insufficient to recommend for or against routinely providing the intervention. The EBPP makes no recommendation. Evidence that the intervention is effective is lacking, of poor quality, or conflicting and the balance of benefits, harms, and costs cannot be determined. |
| Insufficient – NOT Recommended (Consensus-based) | I | The evidence is insufficient for an evidence-based recommendation. The intervention is not recommended for appropriate patients because of high costs/high potential for harm to the patient. |
| NOT Recommended | C | Recommendation against routinely providing the intervention. The EBPP found at least moderate evidence that harms and costs exceed benefits based on limited evidence. |
| Moderately NOT Recommended | B | Recommendation against routinely providing the intervention to eligible patients. The EBPP found at least moderate evidence that the intervention is ineffective, or that harms or costs outweigh benefits. |

| Recommendation Category | Evidence Rating | Description of Category |
|---------------------------------|-----------------|---|
| Strongly NOT Recommended | A | Strong recommendation against providing the intervention to eligible patients. The EBPP found high quality evidence that the intervention is ineffective, or that harms or costs outweigh benefits. |

* "Appropriate" means meeting screening or preventive method entry criteria without contraindications, or having the appropriate diagnosis, indication, time frame, prior conservative testing or treatment, and lack of contraindications for the specific test or treatment.

** For example, would include acetaminophen, and self-administered cold or heat treatments. Excludes all interventional treatments, manual adjustment, and prescriptions medications. Aggregate and individual harms and costs are considered.

COST ANALYSIS

Published cost analyses were reviewed.

METHOD OF GUIDELINE VALIDATION

External Peer Review
Internal Peer Review

DESCRIPTION OF METHOD OF GUIDELINE VALIDATION

Following the chapter and literature review, participants provided written or verbal comments to the American College of Occupational and Environmental Medicine's Practice Guidelines Committee.

Verbal comments were in the form of participation in multi-specialty conference calls, during which the issues raised in each chapter were extensively discussed. Draft chapters were prepared and distributed by the American College of Occupational and Environmental Medicine to all chapter reviewers. Follow-up multi-specialty teleconferences were then held as appropriate, during which time the draft was again reviewed.

2007 Update

Specialty society and society representatives (listed in the original guideline document) served as external reviewers of the low back chapter.

RECOMMENDATIONS

MAJOR RECOMMENDATIONS

Definitions for the strength of evidence ratings (A, B, C, and I) and the criteria for evidence-based recommendations are presented at the end of the "Major Recommendations" field.

General Summary of Recommendations

- The initial assessment of patients with low back problems focuses on detecting indications of potentially serious disease, termed "red flags" (i.e., fever or major trauma).
- In the absence of red flags, imaging and other tests are not recommended in the first 4 to 6 weeks of low back symptoms as they almost never result in a meaningful change in clinical management. Nonprescription medication or an appropriately selected nonsteroidal anti-inflammatory drug (NSAID), appropriate adjustment of physical activity if needed, and the use of thermal modalities such as heat and/or cryotherapies can safely relieve discomfort.
- In the absence of red flags, primary care and occupational physicians or other health care professionals can effectively manage low back problems conservatively.
- At the first visit, the physician should assure the patient that low back pain (LBP) is normal, has an excellent prognosis and, in most cases, is not debilitating on a long-term basis. Patients with elevated fear avoidance beliefs may require additional instructions and interventions to be reassured of this prognosis. Theoretically, this reassurance has the potential to avoid increasing the probability of the patient developing chronic pain syndrome.
- To avoid undue back irritation and debilitation from inactivity, some activity or job modification may be helpful in the acute period. However, bed rest is not recommended for essentially all LBP and radiculopathy patients other than those with unstable fractures or cauda equina syndrome with pending neurological catastrophe. Maintaining ordinary activity, as tolerated, leads to the most rapid recovery.
- All patients should be encouraged to return to work as soon as possible as evidence suggests this leads to the best outcomes. This process may be facilitated with modified duty particularly if job demands exceed patient capabilities. Full-duty work is a reasonable option for patients with low physical job demands and the ability to control such demands (e.g., alternate their posture) as well as for those with less severe presentations.
- Aerobic exercise has the best evidence of efficacy among the exercise regimens, whether for acute, subacute, or chronic LBP patients.
- Non-specific stretching is not recommended as it is not helpful for treatment of LBP. However, specific types of stretching exercises appear helpful (e.g., directional and slump stretching). Strengthening exercises, including lumbar stabilization exercises, are recommended, but not until the acute period of LBP has subsided.
- There is evidence of efficacy for manipulation for treatment of non-specific LBP, particularly for those patients who test positive for the Clinical Prediction Rule.
- Many invasive and noninvasive therapies are intended to cure or manage LBP, but no strong evidence exists that they accomplish this as successfully as therapies that focus on restoring functional ability without focusing on pain. In those cases, the traditional medical model of "curing" the patient does not work well. Furthermore, patients should be aware that returning to normal activities most often aids functional recovery.
- Patients should be encouraged to accept responsibility for managing their recovery rather than expecting the provider to provide an easy "cure." This process will promote using activity rather than pain as a guide, and it will make the treatment goal of return to occupational and non-occupational activities more obvious.

- If symptoms persist without improvement, further evaluation is recommended.
- Within the first 3 months of low back symptoms, only patients with evidence of severe spinal disease or severe debilitating symptoms and physiologic evidence of specific nerve root compromise confirmed by appropriate imaging studies, can be expected to potentially benefit from surgery.
- Quality evidence exists indicating that patient outcomes are not adversely affected by delaying surgery for weeks or a few months and continued conservative care is encouraged in patients with stable or improving deficits who desire to avoid surgery. However, patients with severe or progressive deficits that are not improving at 4 to 6 weeks may benefit from earlier surgical intervention.
- Nonphysical factors (such as psychiatric, psychosocial, workplace, or socioeconomic problems) should be investigated and addressed in cases of delayed recovery or delayed return to work.
- Physicians can greatly improve patient response to back symptoms by providing assurance, encouraging activity, and emphasizing that more than 90% of LBP complaints resolve without any specific therapies. While patients may be looking for a clear-cut diagnosis for their LBP, the risk to them of a suggested "cure" for this assumed diagnosis, resulting in failed expectations, may be worse than their symptoms.
- Physicians should be aware that "abnormal" findings on x-rays, magnetic resonance images, and other diagnostic tests are so common they *are normal* by age 40. Bulging discs continue to increase after age 40, and by age 60 will be encountered in 80% of patients. This requires that a careful history and physical examination be conducted by a skilled physician in order to correlate historical, clinical, and imaging findings prior to assigning the finding on imaging to a patient's complaints. It is recommended that physicians unable to make those correlations, and thus properly educate patients about these complex issues, should defer ordering imaging studies to a qualified consultant in musculoskeletal disorders. Without proper education on prevalence, treatment, and prognosis, patients may become fixated on "fixing" their abnormality (which may in fact be a completely normal condition) and thus iatrogenically increase their risk of developing chronic pain.
- Significant abnormalities in hip range-of-motion may increase the probability of back disorders.

Summary Tables: Recommendations and Evidence

The following summary tables contain the recommendations of the Evidence-based Practice Spine Panel. These recommendations are based on critically appraised higher quality research evidence or, when higher quality evidence was unavailable or inconsistent, on expert consensus observing the First Principles of Clinical Logic as required in the American College of Occupational and Environmental Medicine (ACOEM) Methodology. Table 1 is a summary of the recommendations by treatment (i.e., medications). Table 2 is a summary by low back disorder. The reader is cautioned to utilize the more detailed indications, specific appropriate diagnoses, temporal sequencing, preceding testing or conservative treatment, and contraindications that are elaborated in more detail for each test or treatment in the body of this Guideline when using these

recommendations in clinical practice or medical management. These recommendations are not simple "yes/no" criteria.

Recommendations are made under the following categories:

- Strongly Recommended, "A" Level
- Moderately Recommended, "B" Level
- Recommended, "C" Level
- Insufficient – Recommended (Consensus-based), "I" Level
- Insufficient – No Recommendation (Consensus-based), "I" Level
- Insufficient – Not Recommended (Consensus-based), "I" Level
- Not Recommended, "C" Level
- Moderately Not Recommended, "B" Level
- Strongly Not Recommended, "A" Level

Table 1: Summary of Recommendations for Evaluating and Managing Low Back Disorders

| Clinical Measure | Treatment with Evidence Rating/Recommendation Level | | |
|---------------------------|---|--|--|
| | Recommended | No Recommendation | Not Recommended |
| Diagnostic Testing | <p>X-rays for acute low back pain (LBP) with red flags for fractures or systemic illness, subacute not improving, or chronic LBP as an option to rule out other conditions (I)</p> <p>Flexion and extension views for evaluation of symptomatic spondylolisthesis (I)</p> <p>Magnetic resonance imaging (MRI) for acute LBP during first 6 weeks if red flags (I)</p> <p>MRI for subacute or chronic radicular pain syndromes lasting at least 4 to 6 weeks (B)</p> | <p>Functional capacity evaluations for subacute or chronic stable LBP or post-operative recovery (I)</p> | <p>Routine x-rays for acute, nonspecific LBP (C)</p> <p>MRI for acute radicular pain syndromes in first 6 weeks, regardless of signs of neurological impingement, unless severe and not trending towards improvement (C)</p> <p>Standing or weight-bearing MRI for any back or radicular pain syndrome or condition (I)</p> <p>Computerized tomography (CT) for acute, subacute, chronic non-specific LBP, or radicular pain syndromes (I)</p> <p>Electrodiagnostic study for acute, subacute or chronic LBP patients who do not have significant leg pain or numbness (C)</p> |

| Clinical Measure | Treatment with Evidence Rating/Recommendation Level | | |
|------------------|--|-------------------|--|
| | Recommended | No Recommendation | Not Recommended |
| | <p>MRI as an option for select chronic LBP (I)</p> <p>Computerized tomography (CT) for acute or subacute radicular pain syndrome that has failed to improve within 4 to 6 weeks (C)</p> <p>Myelography, including CT myelography, for uncommon specific situations (I)</p> <p>Electrodiagnostic studies, which must include needle electromyography (EMG) where CT or MRI is equivocal and there are ongoing pain complaints (C)</p> | | <p>Routine bone scanning for LBP (I)</p> <p>Single proton emission computed tomography (SPECT) for acute, subacute, or chronic LBP, or radicular pain syndromes or other LBP-related conditions (I)</p> <p>Diagnostic ultrasound (I)</p> <p>Fluoroscopy for acute, subacute, or chronic LBP (I)</p> <p>Videofluoroscopy for acute, subacute, or chronic LBP (I)</p> <p>Discography for acute, subacute, or chronic LBP or radicular pain syndromes (B)</p> <p>MRI discography (C)</p> <p>Myeloscopy for acute, subacute, or chronic LBP, or spinal stenosis, radicular pain syndromes, or post-surgical back pain problems (I)</p> <p>Surface electromyography (I)</p> <p>Thermography for acute, subacute, or chronic LBP or radicular pain (I)</p> <p>Functional capacity evaluations for acute LBP, acute or subacute radicular pain syndromes, or post-surgical back pain within first 12 weeks of</p> |

| Clinical Measure | Treatment with Evidence Rating/Recommendation Level | | |
|--------------------|---|---|--|
| | Recommended | No Recommendation | Not Recommended |
| | | | postoperative period (I) |
| Medications | <p>NSAIDs for acute LBP (A)</p> <p>NSAIDs for subacute, chronic, or post-operative LBP (B)</p> <p>NSAIDs for radicular pain syndromes including sciatica (C)</p> <p>Cytoprotective medications for patients with contraindications for NSAIDs (C)</p> <p>Acetaminophen for LBP with or without radicular symptoms if contraindications for NSAIDs (C)</p> <p>Discuss risks/benefits of NSAID therapy with patients with known or multiple risk factors for cardiovascular disease (I)</p> <p>Acetaminophen or aspirin as 1st-line therapy for patients with known or multiple risk factors for cardiovascular disease (A)</p> <p>Norepinephrine reuptake inhibitors for chronic LBP (A)</p> | <p>Gabapentin for chronic radicular pain syndromes (I)</p> <p>Thiocolchicoside for acute, subacute, or chronic LBP (I)</p> <p>Creams and ointments for acute, subacute, chronic LBP (I)</p> <p>Camphora molmol, Maleluca alternifolia, Angelica sinensis, Aloe vera, Thymus officinalis, Menthe piperita, Arnica Montana, Curcuma longa, Tanacetum parthenium, Zingiber officinalis (I)</p> | <p>Selective serotonin reuptake inhibitors (e.g., paroxetine, bupropion, trazodone) for chronic LBP (A)</p> <p>Antidepressants for acute or subacute LBP (I)</p> <p>Topiramate for neuropathic pain, including peripheral neuropathy (I)</p> <p>Oral and intravenous (IV) colchicine for acute, subacute, or chronic LBP (I)</p> <p>Gabapentin for chronic non-neuropathic pain or LBP (C)</p> <p>Routine use of opioids for acute, subacute, or chronic LBP (C)</p> <p>Skeletal muscle relaxants for mild to moderate acute LBP or chronic use in subacute or chronic LBP (other than acute exacerbations) (I)</p> <p>Glucocorticosteroids for acute LBP (B)</p> <p>Glucocorticosteroids for subacute or chronic LBP, mild to moderate radiculopathy (I)</p> <p>Tumor necrosis factor-alpha inhibitors for radicular pain syndromes (C)</p> |

| Clinical Measure | Treatment with Evidence Rating/Recommendation Level | | |
|------------------|---|-------------------|--|
| | Recommended | No Recommendation | Not Recommended |
| | <p>Norepinephrine reuptake inhibitors for radicular pain (C)</p> <p>Topiramate for limited use in select chronic LBP as 4th- or 5th-line agent (C)</p> <p>Carbamazepine for chronic radicular or neuropathic pain as a 4th- or 5th-line agent (I)</p> <p>Gabapentin for perioperative pain management (A)</p> <p>Gabapentin for severe neurogenic claudication with limited walking distance (C)</p> <p>Limited use (2 to 3 weeks) of opioids with longer periods for more invasive procedures (C)</p> <p>Skeletal muscle relaxants as 2nd-line treatment in moderate to severe acute LBP not adequately controlled by NSAIDs (B)</p> <p>Skeletal muscle relaxants as 2nd- or 3rd-line treatments for acute radicular pain syndromes or acute post-surgical</p> | | <p>Tumor necrosis factor-alpha inhibitors for acute, subacute, or chronic LBP (I)</p> <p>Vitamins for acute, subacute, or chronic LBP, or post-operative LBP or radiculopathy (I)</p> <p>Willow bark (salix) (I)</p> <p>Spiroflor (I)</p> <p>Complementary or alternative treatments or dietary supplements, etc., other than those discussed in chapter (I)</p> |

| Clinical Measure | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------------------------|---|--|---|
| | Recommended | No Recommendation | Not Recommended |
| | <p>situations (I)</p> <p>Glucocorticosteroids for acute severe radicular pain syndromes (C)</p> <p>Harpagoside in carefully selected patients for acute, subacute, or chronic LBP if NSAIDs contraindicated (C)</p> <p>Capsicum for acute and subacute LBP, or temporary flare-ups of chronic LBP (B)</p> | | |
| Orthotics and Immobilization | <p>Bed rest for unstable spinal fractures (I)</p> <p>Alteration of sleep posture (I)</p> | | <p>Bed rest for acute LBP (A)</p> <p>Bed rest for subacute and chronic LBP (B)</p> <p>Bed rest for stable spinal fractures (I)</p> <p>Bed rest for radicular pain syndromes including sciatica (C)</p> <p>Bed rest for other low back problems (I)</p> <p>Commercial sleeping products (e.g., pillows) for primary prevention or treatment of acute, subacute, or chronic LBP (I)</p> |
| Physical Treatment Methods | <p>Shoe lifts for chronic or recurrent LBP with leg length discrepancy of >2cm (I)</p> <p>Shoe insoles for</p> | <p>Shoe insoles for spinal pain patients, including those without prolonged walking requirements (I)</p> | <p>Shoe insoles and lifts for acute LBP (I)</p> <p>Shoe insoles and lifts for subacute or chronic LBP, radicular pain syndromes or other back-related</p> |

| Clinical Measure | Treatment with Evidence Rating/Recommendation Level | | |
|------------------|---|--|---|
| | Recommended | No Recommendation | Not Recommended |
| | <p>chronic LBP with prolonged walking requirements (C)</p> <p>Self-application of low-tech cryotherapies for acute LBP (I)</p> <p>Self-application of heat therapy including a heat wrap (C)</p> <p>Massage for time-limited use in subacute and chronic LBP patients without underlying serious pathology and as an adjunct to a conditioning program with both graded aerobic exercise and strengthening exercises (C)</p> <p>Massage for acute LBP and chronic radicular pain syndromes (I)</p> <p>Transcutaneous electrical neurostimulation (TENS) (single or dual channel) for select use in chronic LBP or chronic radicular pain syndrome as an adjunct for more efficacious treatments (C)</p> <p>Manipulation or mobilization for</p> | <p>Mattress firmness (I)</p> <p>Other optimal sleeping surfaces (e.g., bedding, water beds, hammocks) (I)</p> <p>Infrared therapy for acute LBP (I)</p> <p>Infrared therapy for home use (I)</p> <p>Ultrasound (I)</p> <p>Neuroreflexotherapy for acute or subacute LBP or radicular pain syndromes (I)</p> <p>Interferential therapy for acute LBP with or without radicular pain (I)</p> | <p>conditions other than leg length discrepancy >2cm (I)</p> <p>Shoe insoles and lifts for primary prevention (C)</p> <p>Routine use of cryotherapies in health care provider offices or home use of a high-tech device for LBP (I)</p> <p>Lumbar supports (C)</p> <p>Lumbar supports for prevention of LBP (C)</p> <p>Magnets (I)</p> <p>Diathermy for any LBP-related condition (C)</p> <p>Infrared therapy for subacute and chronic LBP (I)</p> <p>Low-level laser therapy (I)</p> <p>Mechanical devices for administering massage (C)</p> <p>Traction for acute, subacute, or chronic LBP or radicular pain syndromes (C)</p> <p>Decompression through traction and spinal decompressive devices for acute, subacute, chronic LBP or radicular pain syndromes (I)</p> <p>Interferential therapy for subacute or chronic LBP,</p> |

| Clinical Measure | Treatment with Evidence Rating/Recommendation Level | | |
|------------------|--|-------------------|--|
| | Recommended | No Recommendation | Not Recommended |
| | <p>select acute LBP based on Clinical Prediction Rule (B)</p> <p>Manipulation or mobilization for acute or subacute LBP without Clinical Prediction Rule (C)</p> <p>Acupuncture for select use in chronic LBP as a limited course during which time there are clear objective and functional goals (C)</p> <p>Neuroreflexotherapy for moderate to severe chronic LBP in patients who have failed management with NSAIDs, progressive aerobic exercise program or other exercises, and manipulation (C)</p> | | <p>chronic radicular pain syndromes, or other back-related conditions (C)</p> <p>TENS for acute or subacute LBP or acute radicular pain syndromes (I)</p> <p>Percutaneous electrical nerve stimulation (PENS) for acute or subacute LBP, radicular pain syndromes (I)</p> <p>PENS for chronic non-radicular LBP (I)</p> <p>Microcurrent electrical stimulation for acute, subacute, or chronic LBP or radicular pain syndrome (I)</p> <p>H-wave stimulation for acute, subacute, or chronic LBP or radicular pain syndromes (I)</p> <p>Taping or kinesiotaping for acute, subacute, or chronic LBP, radicular pain syndromes or other back-related conditions (I)</p> <p>Myofascial release for acute, subacute, or chronic LBP, or radicular pain syndromes or other back-related conditions (I)</p> <p>High-voltage galvanic for acute, subacute, or chronic LBP, or radicular pain syndromes or other</p> |

| Clinical Measure | Treatment with Evidence Rating/Recommendation Level | | |
|------------------------------|--|------------------------------------|---|
| | Recommended | No Recommendation | Not Recommended |
| | | | <p>back-related conditions (I)</p> <p>Iontophoresis for acute, subacute, or chronic LBP, or radicular pain syndromes or other back-related conditions (I)</p> <p>Regular or routine manipulation or mobilization (several times a month for years) (I)</p> <p>Manipulation for radicular pain syndromes with acute neurological deficits (I)</p> <p>Adjustments/manipulation of neck/cervical spine, or areas outside lumbopelvic region (I)</p> <p>Manipulation under anesthesia (MUA) and medication-assisted spinal manipulation (MASM) for acute, subacute, or chronic LBP (I)</p> <p>Acupuncture for acute or subacute LBP, radicular pain syndromes or conditions other than chronic LBP (I)</p> <p>Reflexology for chronic LBP (C)</p> <p>Reflexology for acute, or subacute LBP, or other spinal conditions (I)</p> |
| Activity and Exercise | Aerobic exercise for acute, subacute, or chronic LBP (A) | Yoga for acute or subacute LBP (I) | Aggressive stretching exercises (I) |

| Clinical Measure | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------|---|---|--|
| | Recommended | No Recommendation | Not Recommended |
| | <p>Aerobic exercise for post-operative patients (I)</p> <p>Slump stretch-related exercise or directional preference stretching exercises for acute, subacute, or chronic LBP (C)</p> <p>Strengthening exercises (after instituting aerobic exercises) for acute, subacute, or chronic LBP, post-operative LBP patients (C)</p> <p>Inclusion of Fear Avoidance Belief Training (FABT) during course of rehabilitation (I)</p> <p>Yoga for select highly motivated chronic LBP patients (C)</p> <p>Trial of aquatic therapy for subacute or chronic LBP if patient meets referral criteria for supervised exercise therapy and has co-morbidities that preclude participation in weight-bearing physical activity (I)</p> | | <p>Stretching exercises for preventing LBP (C)</p> <p>Strengthening of abdominal muscles for treatment or prevention of LBP (I)</p> <p>Lumbar extension machines for acute, subacute, or chronic LBP, or any radicular pain syndrome (I)</p> <p>Aquatic therapy for all acute LBP and all other subacute and chronic LBP not meeting referral criteria (I)</p> |
| Injections | Epidural glucocorticosteroid injections as option | Diagnostic facet joint injections for chronic LBP (I) | Epidural glucocorticosteroid injections for acute, |

| Clinical Measure | Treatment with Evidence Rating/Recommendation Level | | |
|------------------|---|---|---|
| | Recommended | No Recommendation | Not Recommended |
| | <p>for acute or subacute radicular pain syndromes lasting at least 3 weeks after treating with NSAIDs and without evidence of trending towards spontaneous resolution (I)</p> <p>Epidural glucocorticosteroid injections as 2nd-line treatment of acute spinal stenosis flare-ups (I)</p> <p>Trigger and/or tender point injections as 2nd or 3rd option for subacute or chronic LBP that is not resolving (C)</p> <p>Sacroiliac joint corticosteroid injections as option for patients with specific known cause of sacroiliitis (C)</p> | <p>Botulinum injections for acute, subacute, or chronic LBP, or radicular pain syndromes or other low back problems (I)</p> | <p>subacute, or chronic LBP in the absence of radicular signs and symptoms (C)</p> <p>Intradiscal steroids for acute LBP (I)</p> <p>Intradiscal steroids for subacute or chronic LBP (B)</p> <p>Trigger and/or tender point injections for acute LBP (I)</p> <p>Diagnostic facet joint injections for acute or subacute LBP or radicular pain syndromes (I)</p> <p>Therapeutic facet joint injections for acute, subacute, chronic LBP or radicular pain syndrome (B)</p> <p>Facet joint hyaluronic acid injections (I)</p> <p>Sacroiliac joint injections for acute LBP, including LBP thought to be sacroiliac joint related (I)</p> <p>Prolotherapy injections for acute, subacute, chronic LBP or radicular pain syndrome (C)</p> <p>Radiofrequency lesioning of dorsal root ganglia for chronic sciatica (B)</p> <p>Radiofrequency neurotomy, neurotomy, or facet rhizotomy for any spinal condition (C)</p> |

| Clinical Measure | Treatment with Evidence Rating/Recommendation Level | | |
|--------------------------------|---|-------------------|---|
| | Recommended | No Recommendation | Not Recommended |
| | | | <p>Intradiscal electrothermal annuloplasty (IDET) for acute, subacute, or chronic LBP or any other back-related disorder (I)</p> <p>Percutaneous intradiscal radiofrequency thermocoagulation (PIRFT) for acute, subacute, or chronic LBP, including discogenic LBP (A)</p> |
| Surgical Considerations | <p>Lumbar discectomy for radiculopathy due to ongoing nerve root compression with continued significant pain and functional limitation after 4 to 6 weeks and appropriate conservative treatment (B)</p> <p>Decompressive surgery for symptomatic spinal stenosis (neurogenic claudication) that is intractable to conservative management (B)</p> <p>Lumbar fusion for isthmic spondylolisthesis (C)</p> <p>Lumbar fusion for degenerative spondylolisthesis (C)</p> | | <p>Percutaneous discectomy (nucleoplasty), laser discectomy, and disc coblation therapy for any back or radicular pain syndrome (B)</p> <p>Discectomy for acute, subacute, or chronic LBP without radiculopathy (B)</p> <p>Lumbar fusion for spinal stenosis unless concomitant instability or deformity proven (C)</p> <p>Lumbar fusion for radiculopathy from disc herniation or chronic LBP after lumbar discectomy (C)</p> <p>Lumbar fusion for chronic non-specific LBP (B)</p> <p>Artificial disc replacement for chronic non-specific LBP or other spinal pain syndrome (I)</p> <p>Sacroiliac joint fusion surgery and other sacroiliac joint surgical</p> |

| Clinical Measure | Treatment with Evidence Rating/Recommendation Level | | |
|--|---|-------------------|---|
| | Recommended | No Recommendation | Not Recommended |
| | <p>For 3rd lumbar discectomy on same disc, spine fusion at time of discectomy is an option (I)</p> <p>Vertebroplasty and kyphoplasty for select patients (I)</p> | | <p>procedures for any LBP condition (I)</p> <p>Spinal cord stimulators for acute, subacute, or chronic LBP, or radicular pain syndromes or failed back surgery syndrome (I)</p> <p>Adhesiolysis for acute, subacute, or chronic LBP, or spinal stenosis, or radicular pain syndromes (I)</p> |
| Rehabilitation/ Behavioral/ Education | <p>Chronic pain management or functional restoration program for chronic pain management (I)</p> <p>Chronic pain management or functional restoration program for subacute LBP (I)</p> <p>Work conditioning and work hardening programs for chronic LBP (C)</p> <p>Work conditioning and work hardening programs for subacute LBP (I)</p> <p>Participatory ergonomic programs for highly select subacute and chronic LBP (C)</p> <p>Biofeedback for select chronic LBP as component of an</p> | | <p>Back school for acute LBP (I)</p> <p>Back school and education for prevention of LBP (C)</p> <p>Cognitive behavioral therapy for acute LBP (I)</p> <p>Chronic pain management or functional restoration program for acute spinal disorders (I)</p> <p>Work conditioning and work hardening programs for acute LBP (I)</p> <p>Biofeedback for acute or subacute LBP (I)</p> <p>Multidisciplinary rehabilitation program with primary focus on LBP interventions (I)</p> |

| Clinical Measure | Treatment with Evidence Rating/Recommendation Level | | |
|------------------|--|-------------------|-----------------|
| | Recommended | No Recommendation | Not Recommended |
| | <p>interdisciplinary approach (I)</p> <p>Multidisciplinary rehabilitation programs with focus on cognitive behavioral, occupational, and activity-based approaches combined with aerobic exercise and other conditioning exercise for chronic LBP (C)</p> <p>Multidisciplinary rehabilitation program with participatory ergonomics team for subacute or chronic LBP with lost-time injuries (C)</p> <p>Smoking cessation and weight loss programs to prevent LBP (I)</p> <p>FABT for acute, subacute, or chronic LBP (B)</p> <p>Back school and education for select chronic LBP and chronic radicular pain syndromes (B)</p> <p>Cognitive behavioral therapy as component of interdisciplinary program for chronic</p> | | |

| Clinical Measure | Treatment with Evidence Rating/Recommendation Level | | |
|------------------|--|-------------------|-----------------|
| | Recommended | No Recommendation | Not Recommended |
| | LBP and subacute LBP when combined with other indicated therapies with parameters described in "Rehabilitation for Delayed Recovery" section (C) | | |

Table 2: Summary of Recommendations by Low Back Disorders

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|----------------------------|--|--|---|
| | Recommended | No Recommendation | Not Recommended |
| Acute Low Back Pain | <p>NSAIDs (A)</p> <p>Cytoprotective medications particularly if contraindications for NSAIDs (C)</p> <p>Acetaminophen if contraindications for NSAIDs (C)</p> <p>Discuss risks/benefits of NSAID therapy with patients with known or multiple risk factors for cardiovascular disease (I)</p> <p>Acetaminophen or aspirin as 1st-line therapy for patients with known or multiple risk factors for cardiovascular disease (A)</p> <p>Limited use (2 to 3 weeks) of opioids</p> | <p>Thiocolchicoside (I)</p> <p>Creams and ointments (I)</p> <p>Camphora molmol, Maleluca alternifolia, Angelica sinensis, Aloe vera, Thymus officinalis, Menthe piperita, Arnica Montana, Curcuma longa, Tanacetum parthenium, Zingiber officinalis (I)</p> <p>Mattress firmness (I)</p> <p>Other optimal sleeping surfaces (e.g., bedding, water beds, hammocks) (I)</p> <p>Infrared therapy (I)</p> <p>Infrared therapy for home use (I)</p> | <p>Antidepressants (I)</p> <p>Anti-epileptic agents including carbamazepine (I)</p> <p>Oral and IV colchicine (I)</p> <p>Routine use of opioids (C)</p> <p>Skeletal muscle relaxants for mild to moderate acute LBP pain (I)</p> <p>Glucocorticosteroids (B)</p> <p>Tumor necrosis factor-alpha inhibitors (I)</p> <p>Vitamins (I)</p> <p>Willow bark (salix) (I)</p> <p>Spiroflor (I)</p> <p>Complementary or alternative treatments or dietary supplements, etc., other than those discussed in chapter (I)</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------|--|---|--|
| | Recommended | No Recommendation | Not Recommended |
| | <p>with longer periods for more invasive procedures (C)</p> <p>Skeletal muscle relaxants as 2nd-line treatment in select cases of moderate to severe acute LBP (B)</p> <p>Harpagoside in carefully selected patients if NSAIDs contraindicated (C)</p> <p>Capsicum (B)</p> <p>Alteration of sleep posture (I)</p> <p>Self-application of low-tech cryotherapies (I)</p> <p>Self-application of heat therapy including a heat wrap (C)</p> <p>Massage (I)</p> <p>Manipulation or mobilization for select patients based on Clinical Prediction Rule (B)</p> <p>Manipulation or mobilization for LBP without Clinical Prediction Rule (C)</p> <p>Aerobic exercise (A)</p> <p>Slump stretch-related exercise or directional</p> | <p>Interferential therapy—with or without radicular pain (I)</p> <p>Ultrasound (I)</p> <p>Neuroreflexotherapy (I)</p> <p>Yoga (I)</p> <p>Botulinum injections (I)</p> | <p>Bed rest (A)</p> <p>Commercial sleeping products for primary prevention or treatment (I)</p> <p>Shoe insoles and lifts (I)</p> <p>Reflexology (I)</p> <p>Lumbar supports (C)</p> <p>Magnets (I)</p> <p>Routine use of cryotherapies in health care provider offices or home use of high-tech device (I)</p> <p>Diathermy (C)</p> <p>Low-level laser therapy (I)</p> <p>Mechanical devices for administering massage (C)</p> <p>Traction (C)</p> <p>Decompression through traction and spinal decompressive devices (I)</p> <p>TENS (I)</p> <p>PENS (I)</p> <p>Microcurrent electrical stimulation (I)</p> <p>H-wave stimulation (I)</p> <p>Taping and kinesiotaping (I)</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------|--|-------------------|---|
| | Recommended | No Recommendation | Not Recommended |
| | <p>preference stretching exercises (C)</p> <p>Strengthening exercises after aerobic exercises instituted (C)</p> <p>Fear Avoidance Belief Training (FABT) (B)</p> <p>Inclusion of FABT during course of rehabilitation (I)</p> | | <p>Myofascial release (I)</p> <p>High-voltage galvanic (I)</p> <p>Iontophoresis (I)</p> <p>Adjustments/manipulation of neck/cervical spine or areas outside lumbopelvic region (I)</p> <p>Manipulation under anesthesia (MUA) and medication-assisted spinal manipulation (MASM) (I)</p> <p>Acupuncture (I)</p> <p>Strengthening of abdominal muscles (I)</p> <p>Aggressive stretching (I)</p> <p>Aquatic therapy (I)</p> <p>Lumbar extension machines (I)</p> <p>Epidural glucocorticosteroid injections in absence of radicular signs and symptoms (C)</p> <p>Intradiscal steroids (I)</p> <p>Trigger and/or tender point injections (I)</p> <p>Diagnostic facet joint injections (I)</p> <p>Therapeutic facet joint injections (B)</p> <p>Facet joint hyaluronic acid injections (I)</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------------------|--|--|--|
| | Recommended | No Recommendation | Not Recommended |
| | | | <p>Sacroiliac joint injections (I)</p> <p>Prolotherapy injections (C)</p> <p>Radiofrequency neurotomy, neurotomy, or facet rhizotomy (C)</p> <p>IDET (I)</p> <p>PIRFT particularly including discogenic LBP (A)</p> <p>Spinal cord stimulators (I)</p> <p>Discectomy for acute LBP without radiculopathy treatment (B)</p> <p>Adhesiolysis (I)</p> <p>Back school (I)</p> <p>Cognitive behavioral therapy (I)</p> <p>Chronic pain management or functional restoration program (I)</p> <p>Work conditioning and work hardening programs (I)</p> <p>Biofeedback (I)</p> |
| Subacute Low Back Pain | <p>NSAIDs (B)</p> <p>Cytoprotective medications particularly if contraindications for NSAIDs (C)</p> <p>Acetaminophen if</p> | <p>Thiocolchicoside (I)</p> <p>Creams and ointments (I)</p> <p>Camphora molmol, Maleluca alternifolia, Angelica sinensis, Aloe vera,</p> | <p>Antidepressants (I)</p> <p>Anti-epileptic agents including carbamazepine (I)</p> <p>Oral and IV colchicine (I)</p> <p>Gabapentin (I)</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------|--|--|--|
| | Recommended | No Recommendation | Not Recommended |
| | <p>contraindications for NSAIDs (C)</p> <p>Discuss risks/benefits of NSAID therapy with patients with known or multiple risk factors for cardiovascular disease (I)</p> <p>Acetaminophen or aspirin as 1st-line therapy for patients with known or multiple risk factors for cardiovascular disease (A)</p> <p>Harpagoside in carefully selected patients if NSAIDs contraindicated (C)</p> <p>Capsicum (B)</p> <p>Alteration of sleep posture (I)</p> <p>Self-application of low-tech cryotherapies (I)</p> <p>Self-application of heat therapy including a heat wrap (C)</p> <p>Massage for time limited use in subacute LBP patients without underlying serious pathology and as an adjunct to a conditioning program that has</p> | <p>Thymus officinalis, Menthe piperita, Arnica Montana, Curcuma longa, Tanacetum parthenium, Zingiber officinalis (I)</p> <p>Mattress firmness (I)</p> <p>Other optimal sleeping surfaces, e.g., bedding, water beds, hammocks (I)</p> <p>Infrared therapy for home use (I)</p> <p>Ultrasound (I)</p> <p>Neuroreflexotherapy (I)</p> <p>Yoga (I)</p> <p>Botulinum injections (I)</p> | <p>Routine use of opioids (C)</p> <p>Skeletal muscle relaxants (I)</p> <p>Glucocorticosteroids (I)</p> <p>Tumor necrosis factor-alpha inhibitors (I)</p> <p>Vitamins (I)</p> <p>Willow bark (salix) (I)</p> <p>Spiroflor (I)</p> <p>Complementary or alternative treatments or dietary supplements, etc., other than those discussed in chapter (I)</p> <p>Bed rest (B)</p> <p>Commercial sleeping products for primary prevention or treatment (I)</p> <p>Shoe insoles and lifts except if leg length discrepancy >2cm (I)</p> <p>Reflexology (I)</p> <p>Lumbar supports (C)</p> <p>Magnets (I)</p> <p>Routine use of cryotherapies in health care provider offices or home use of high-tech device (I)</p> <p>Diathermy (C)</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------|--|-------------------|---|
| | Recommended | No Recommendation | Not Recommended |
| | <p>both graded aerobic exercise and strengthening exercises (C)</p> <p>Manipulation or mobilization for LBP without Clinical Prediction Rule (C)</p> <p>Aerobic exercise (A)</p> <p>Slump stretch-related exercise or directional preference stretching exercises (C)</p> <p>Strengthening exercises after aerobic exercises instituted (C)</p> <p>Trial of aquatic therapy if patient meets referral criteria for supervised exercise therapy and has co-morbidities that preclude participation in weight-bearing physical activity (I)</p> <p>Trigger and/or tender point injections as 2nd or 3rd option for subacute LBP that is not resolving (C)</p> <p>Chronic pain management or functional restoration program (I)</p> | | <p>Infrared therapy (I)</p> <p>Low-level laser therapy (I)</p> <p>Mechanical devices for administration of massage (C)</p> <p>Traction (C)</p> <p>Decompression through traction and spinal decompressive devices (I)</p> <p>Interferential therapy (C)</p> <p>TENS (I)</p> <p>PENS (I)</p> <p>Microcurrent electrical stimulation (I)</p> <p>H-wave stimulation (I)</p> <p>Taping and kinesiotaping (I)</p> <p>Myofascial release (I)</p> <p>High-voltage galvanic (I)</p> <p>Iontophoresis (I)</p> <p>Adjustments/manipulation of neck/cervical spine or areas outside lumbopelvic region (I)</p> <p>MUA and MASM (I)</p> <p>Acupuncture (I)</p> <p>Strengthening of abdominal muscles (I)</p> <p>Aggressive stretching (I)</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|--------------------|--|----------------------|--|
| | Recommended | No Recommendation | Not Recommended |
| | <p>Work conditioning and work hardening programs (I)</p> <p>Participatory ergonomic programs for highly select subacute LBP (C)</p> <p>Multidisciplinary rehabilitation program with a participatory ergonomics team for subacute LBP with lost-time injuries (C)</p> <p>Inclusion of FABT during course of rehabilitation (I)</p> <p>FABT (B)</p> <p>Cognitive behavioral therapy as component of interdisciplinary program (C)</p> | | <p>Aquatic therapy for all other subacute LBP (I)</p> <p>Lumbar extension machines (I)</p> <p>Epidural glucocorticosteroid injections in absence of radicular signs and symptoms (C)</p> <p>Intradiscal steroids (B)</p> <p>Diagnostic facet joint injections (I)</p> <p>Therapeutic facet joint injections (B)</p> <p>Facet joint hyaluronic acid injections (I)</p> <p>Prolotherapy injections (C)</p> <p>Radiofrequency neurotomy, neurotomy, facet rhizotomy (C)</p> <p>IDET (I)</p> <p>PIRFT particularly including discogenic LBP (A)</p> <p>Spinal cord stimulators (I)</p> <p>Discectomy for subacute LBP without radiculopathy treatment (B)</p> <p>Adhesiolysis (I)</p> <p>Biofeedback (I)</p> |
| Chronic Low | NSAIDs (B) | Thiocolchicoside (I) | Selective serotonin |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------|--|---|---|
| | Recommended | No Recommendation | Not Recommended |
| Back Pain | <p>Cytoprotective medications if contraindications for NSAIDs (C)</p> <p>Acetaminophen if contraindications for NSAIDs (C)</p> <p>Discuss risks/benefits of NSAID therapy with patients with known or multiple risk factors for cardiovascular disease (I)</p> <p>Acetaminophen or aspirin as the 1st-line therapy for patients with known or multiple risk factors for cardiovascular disease (A)</p> <p>Norepinephrine reuptake inhibitors (A)</p> <p>Opioid trial – both function and pain must improve to continue (I)</p> <p>Topiramate for limited use as 4th- or 5th-line agent (C)</p> <p>Harpagoside in carefully selected patients if NSAIDs contraindicated (C)</p> <p>Capsicum for</p> | <p>Creams and ointments (I)</p> <p>Camphora molmol, Maleluca alternifolia, Angelica sinensis, Aloe vera, Thymus officinalis, Menthe piperita, Arnica Montana, Curcuma longa, Tanacetum parthenium, Zingiber officinalis (I)</p> <p>Mattress firmness (I)</p> <p>Other optimal sleeping surfaces (e.g., bedding, water beds, hammocks) (I)</p> <p>Infrared therapy for home use (I)</p> <p>Ultrasound (I)</p> <p>Diagnostic facet joint injections (I)</p> <p>Botulinum injections (I)</p> | <p>reuptake inhibitors (e.g., paroxetine, bupropion, trazodone) (A)</p> <p>Anti-epileptic agents including carbamazepine (I)</p> <p>Oral and IV colchicine (I)</p> <p>Gabapentin (C)</p> <p>Routine use of opioids (C)</p> <p>Skeletal muscle relaxants (I)</p> <p>Systemic glucocorticosteroids (I)</p> <p>Tumor necrosis factor-alpha inhibitors (I)</p> <p>Vitamins (I)</p> <p>Willow bark (salix) (I)</p> <p>Spiroflor (I)</p> <p>Complementary or alternative treatments or dietary supplements, etc., other than those discussed in chapter (I)</p> <p>Bed rest (B)</p> <p>Commercial sleeping products for primary prevention or treatment (I)</p> <p>Reflexology (C)</p> <p>Shoe insoles and lifts for chronic LBP other than for lg length discrepancy >2cm (I)</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------|---|-------------------|--|
| | Recommended | No Recommendation | Not Recommended |
| | <p>temporary flare-ups (B)</p> <p>Alteration of sleep posture (I)</p> <p>Neuroreflexotherapy for moderate to severe chronic LBP in patients who have failed management with NSAIDs, progressive aerobic exercise program or other exercise and manipulation (C)</p> <p>Shoe insoles if prolonged walking requirements (C)</p> <p>Shoe lifts for chronic or recurrent LBP patients with leg length discrepancy of >2cm (I)</p> <p>Self-application of low-tech cryotherapies (I)</p> <p>Self-application of heat therapy including a heat wrap (C)</p> <p>Massage for time limited use in chronic LBP patients without underlying serious pathology and as an adjunct to a conditioning program that has both graded aerobic exercise and</p> | | <p>Lumbar supports (C)</p> <p>Magnets (I)</p> <p>Routine use of cryotherapies in health care provider offices or home use of high-tech device (I)</p> <p>Diathermy (C)</p> <p>Infrared therapy (I)</p> <p>Low-level laser therapy (I)</p> <p>Mechanical devices for administering massage (C)</p> <p>Traction (C)</p> <p>Decompression through traction and spinal decompressive devices (I)</p> <p>Interferential therapy (C)</p> <p>PENS for chronic nonradicular LBP (I)</p> <p>Microcurrent electrical stimulation (I)</p> <p>H-wave stimulation (I)</p> <p>Taping and kinesiotaping (I)</p> <p>Myofascial release (I)</p> <p>High-voltage galvanic (I)</p> <p>Iontophoresis (I)</p> <p>Regular or routine</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------|---|-------------------|--|
| | Recommended | No Recommendation | Not Recommended |
| | <p>strengthening exercises (C)</p> <p>TENS for chronic LBP as an adjunct for more efficacious treatments (C)</p> <p>Acupuncture for select use as a limited course during which there are clear objective and functional goals (C)</p> <p>Aerobic exercise (A)</p> <p>Slump stretch-related exercise or directional preference stretching exercises (C)</p> <p>Strengthening exercises after aerobic exercises instituted (C)</p> <p>Inclusion of FABT during course of rehabilitation (I)</p> <p>Yoga for select highly motivated patients (C)</p> <p>Trial of aquatic therapy if referral criteria met for supervised exercise therapy and co-morbidities that preclude participation in weight-bearing physical activity (I)</p> | | <p>manipulation or mobilization (several times a month for years) (I)</p> <p>Adjustments/manipulation of neck/cervical spine or areas outside lumbopelvic region (I)</p> <p>MUA and MASM (I)</p> <p>Aggressive stretching exercises (I)</p> <p>Strengthening of abdominal muscles (I)</p> <p>Aquatic therapy for all other chronic LBP (I)</p> <p>Lumbar extension machines (I)</p> <p>Epidural glucocorticosteroid injections in absence of radicular signs and symptoms (C)</p> <p>Intradiscal steroids (B)</p> <p>Therapeutic facet joint injections (B)</p> <p>Facet joint hyaluronic acid injections (I)</p> <p>Prolotherapy injections (C)</p> <p>Radiofrequency neurotomy, neurotomy, facet rhizotomy (C)</p> <p>IDET (I)</p> <p>PIRFT particularly</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------|--|-------------------|---|
| | Recommended | No Recommendation | Not Recommended |
| | <p>Trigger and/or tender point injections as 2nd or 3rd option for chronic LBP that is not resolving (C)</p> <p>Chronic pain management and functional restoration program (I)</p> <p>Work conditioning and work hardening programs (C)</p> <p>Participatory ergonomic programs for highly select chronic LBP (C)</p> <p>Biofeedback for select chronic LBP as component of an interdisciplinary approach (I)</p> <p>Multidisciplinary rehabilitation programs with focus on cognitive behavioral, occupational, and activity-based approaches combined with aerobic exercise and other conditioning exercise (C)</p> <p>Multidisciplinary rehabilitation program with a participatory</p> | | <p>including discogenic LBP (A)</p> <p>Lumbar fusion for chronic LBP after lumbar discectomy (C)</p> <p>Lumbar fusion for chronic nonspecific LBP (B)</p> <p>Artificial disc replacement for chronic nonspecific LBP (I)</p> <p>Spinal cord stimulators (I)</p> <p>Sacroiliac joint fusion surgery and other sacroiliac joint surgical procedures (I)</p> <p>Percutaneous discectomy (nucleoplasty), laser discectomy, and disc coblation therapy (B)</p> <p>Discectomy for chronic LBP without radiculopathy treatment (B)</p> <p>Adhesiolysis (I)</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------------------------|--|--|---|
| | Recommended | No Recommendation | Not Recommended |
| | <p>ergonomics team for chronic LBP with lost-time injuries (C)</p> <p>FABT (B)</p> <p>Back school and education (B)</p> <p>Cognitive behavioral therapy as component of interdisciplinary program (C)</p> | | |
| Post-operative Low Back Pain | <p>NSAIDs (B)</p> <p>Cytoprotective medications if contraindications for NSAIDs (C)</p> <p>Acetaminophen if contraindications for NSAIDs (C)</p> <p>Discuss risks/benefits of NSAID therapy with patients with known or multiple risk factors for cardiovascular disease (I)</p> <p>Acetaminophen or aspirin as 1st-line therapy for patients with known or multiple risk factors for cardiovascular disease (A)</p> <p>Gabapentin for perioperative pain management (A)</p> | <p>Camphora molmol, Maleluca alternifolia, Angelica sinensis, Aloe vera, Thymus officinalis, Menthe piperita, Arnica Montana, Curcuma longa, Tanacetum parthenium, Zingiber officinalis (I)</p> <p>Shoe insoles (I)</p> <p>Mattress firmness (I)</p> <p>Other optimal sleeping surfaces (e.g., bedding, water beds, hammocks) (I)</p> <p>Infrared therapy for home use (I)</p> <p>Ultrasound (I)</p> <p>Botulinum injections (I)</p> | <p>Vitamins (I)</p> <p>Willow bark (salix) (I)</p> <p>Spiroflor (I)</p> <p>Complementary or alternative treatments or dietary supplements, etc., other than those discussed in chapter (I)</p> <p>Bed rest (I)</p> <p>Reflexology (I)</p> <p>Lumbar supports (C)</p> <p>Magnets (I)</p> <p>Routine use of cryotherapies in health care provider offices or home use of high-tech device (I)</p> <p>Diathermy (C)</p> <p>Low-level laser therapy (I)</p> <p>Mechanical devices for</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|---|---|---|---|
| | Recommended | No Recommendation | Not Recommended |
| | <p>Limited use (2 to 3 weeks) of opioids with longer periods for more invasive procedures (C)</p> <p>Skeletal muscle relaxants as 2nd- or 3rd-line agents for acute post-surgical situations (I)</p> <p>Alteration of sleep posture (I)</p> <p>Aerobic exercise (I)</p> <p>Strengthening exercises after aerobic exercises instituted (C)</p> <p>Stretching exercises to regain normal range of motion (I)</p> <p>Inclusion of FABT during course of rehabilitation (I)</p> | | <p>administering massage (C)</p> <p>Interferential therapy (C)</p> <p>Taping and kinesiotaping (I)</p> <p>Myofascial release (I)</p> <p>High-voltage galvanic (I)</p> <p>Iontophoresis (I)</p> <p>Adjustments/manipulation of neck/cervical spine or areas outside lumbopelvic region (I)</p> <p>Acupuncture (I)</p> <p>Aggressive stretching (I)</p> <p>Strengthening of abdominal muscles (I)</p> <p>Radiofrequency neurotomy, neurotomy, facet rhizotomy (C)</p> <p>IDET (I)</p> <p>Spinal cord stimulators for failed back surgery syndrome (I)</p> |
| <p>Radicular Pain Syndromes (including "sciatica")</p> | <p>NSAIDs (C)</p> <p>Cytoprotective medications if contraindications for NSAIDs (C)</p> <p>Acetaminophen if contraindications for NSAIDs (C)</p> <p>Discuss</p> | <p>Camphora molmol, Maleluca alternifolia, Angelica sinensis, Aloe vera, Thymus officinalis, Menthe piperita, Arnica Montana, Curcuma longa, Tanacetum parthenium, Zingiber officinalis (I)</p> | <p>Glucocorticosteroids for mild to moderate radiculopathy (I)</p> <p>Tumor necrosis factor-α inhibitors (C)</p> <p>Willow bark (salix) (I)</p> <p>Spiroflor (I)</p> <p>Vitamins (I)</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------|---|--|---|
| | Recommended | No Recommendation | Not Recommended |
| | <p>risks/benefits of NSAID therapy with patients with known or multiple risk factors for cardiovascular disease (I)</p> <p>Acetaminophen or aspirin as 1st-line therapy for patients with known or multiple risk factors for cardiovascular disease (A)</p> <p>Norepinephrine reuptake inhibitors (e.g., tricyclic antidepressants) (C)</p> <p>Opioid trial – both function and pain must improve to continue (I)</p> <p>Carbamazepine as 4th- or 5th-line treatment (I)</p> <p>Skeletal muscle relaxants as 2nd- or 3rd-line agents for acute radicular pain (I)</p> <p>Glucocorticosteroids for acute, severe radicular pain syndromes (C)</p> <p>Alteration of sleep posture (I)</p> <p>Massage for chronic radicular pain syndromes (I)</p> | <p>Gabapentin for chronic radicular pain syndromes (I)</p> <p>Shoe insoles (I)</p> <p>Mattress firmness (I)</p> <p>Other optimal sleeping surfaces (e.g., bedding, water beds, hammocks) (I)</p> <p>Infrared therapy for home use (I)</p> <p>Ultrasound (I)</p> <p>Neuroreflexotherapy (I)</p> <p>Botulinum injections (I)</p> | <p>Complementary or alternative treatments or dietary supplements, etc., other than those discussed in chapter (I)</p> <p>Bed rest (C)</p> <p>Reflexology (I)</p> <p>Shoe insoles and lifts except if leg length discrepancy >2 cm (I)</p> <p>Lumbar supports (C)</p> <p>Magnets (I)</p> <p>Routine use of cryotherapies in health care provider offices or home use of high-tech device (I)</p> <p>Lumbar extension machines (I)</p> <p>Diathermy (C)</p> <p>Low-level laser therapy (I)</p> <p>Mechanical devices for administering massage (C)</p> <p>Traction (C)</p> <p>Decompression through traction and spinal decompressive devices (I)</p> <p>Interferential therapy for chronic radicular pain (C)</p> <p>TENS for acute radicular pain (I)</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------|--|-------------------|---|
| | Recommended | No Recommendation | Not Recommended |
| | <p>TENS for chronic radicular pain syndrome as an adjunct for more efficacious treatments (C)</p> <p>Epidural glucocorticosteroid injections for acute or subacute radicular pain syndromes lasting at least 3 weeks after treatment with NSAIDs and without evidence of trending towards spontaneous resolution (I)</p> <p>Back school and education for chronic radicular pain syndromes (B)</p> <p>Lumbar discectomy for patients with radiculopathy due to ongoing nerve root compression who continue to have significant pain and functional limitation after 4 to 6 weeks and appropriate conservative treatment (B)</p> <p>For 3rd lumbar discectomy on same disc, spine fusion at time of discectomy is an option (I)</p> | | <p>PENS (I)</p> <p>Microcurrent electrical stimulation (I)</p> <p>H-wave stimulation (I)</p> <p>Taping and kinesiotaping (I)</p> <p>Myofascial release (I)</p> <p>High-voltage galvanic (I)</p> <p>Iontophoresis (I)</p> <p>Manipulation for radicular pain syndromes with acute neurological deficits (I)</p> <p>Adjustments/manipulation of neck/cervical spine or areas outside lumbopelvic region (I)</p> <p>Acupuncture (I)</p> <p>Diagnostic facet joint injections (I)</p> <p>Therapeutic facet joint injections (B)</p> <p>Facet joint hyaluronic acid injections (I)</p> <p>Prolotherapy injections (C)</p> <p>Radiofrequency neurotomy, neurotomy, facet rhizotomy (C)</p> <p>IDET (I)</p> <p>Lumbar fusion for</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|------------------------|--|--|---|
| | Recommended | No Recommendation | Not Recommended |
| | | | <p>radiculopathy from disc herniation (C)</p> <p>Percutaneous discectomy (nucleoplasty), laser discectomy, and disc coblation therapy (B)</p> <p>Spinal cord stimulators (I)</p> <p>Artificial disc replacement (I)</p> <p>Sacroiliac joint fusion surgery and other sacroiliac joint surgical procedures (I)</p> <p>Adhesiolysis (I)</p> |
| Spinal Stenosis | <p>Cytoprotective medications if contraindications for NSAIDs (C)</p> <p>Acetaminophen if contraindications for NSAIDs (C)</p> <p>Discuss risks/benefits of NSAID therapy with patients with known or multiple risk factors for cardiovascular disease (I)</p> <p>Acetaminophen or aspirin as 1st-line therapy for patients with known or multiple risk factors for cardiovascular disease (A)</p> <p>Gabapentin for severe neurogenic</p> | <p>Camphora molmol, Maleluca alternifolia, Angelica sinensis, Aloe vera, Thymus officinalis, Menthe piperita, Arnica Montana, Curcuma longa, Tanacetum parthenium, Zingiber officinalis (I)</p> <p>Shoe insoles (I)</p> <p>Mattress firmness (I)</p> <p>Other optimal sleeping surfaces (e.g., bedding, water beds, hammocks) (I)</p> <p>Infrared therapy for home use (I)</p> <p>Ultrasound (I)</p> | <p>Willow bark (salix) (I)</p> <p>Spiroflor (I)</p> <p>Complementary or alternative treatments or dietary supplements, etc., other than those discussed in chapter (I)</p> <p>Bed rest (I)</p> <p>Shoe insoles and lifts except if leg length discrepancy >2 cm (I)</p> <p>Reflexology (I)</p> <p>Shoe insoles and lifts for primary prevention (C)</p> <p>Lumbar supports (C)</p> <p>Magnets (I)</p> <p>Routine use of cryotherapies in health care provider offices or</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------------|---|---|--|
| | Recommended | No Recommendation | Not Recommended |
| | <p>claudication with limited walking distance from spinal stenosis (C)</p> <p>Opioid trial—both function and pain must improve to continue (I)</p> <p>Alteration of sleep posture (I)</p> <p>Epidural glucocorticosteroid injections as 2nd-line treatment of acute flare-ups (I)</p> <p>Decompressive surgery for symptomatic spinal stenosis that is intractable to conservative management (B)</p> | <p>Botulinum injections (I)</p> | <p>home use of high-tech device (I)</p> <p>Diathermy (C)</p> <p>Low-level laser therapy (I)</p> <p>Mechanical devices for administering massage (C)</p> <p>Interferential therapy (C)</p> <p>Taping and kinesiotaping (I)</p> <p>Myofascial release (I)</p> <p>High voltage galvanic (I)</p> <p>Iontophoresis (I)</p> <p>Adjustments/manipulation of neck/cervical spine or areas outside lumbopelvic region (I)</p> <p>Acupuncture (I)</p> <p>Radiofrequency neurotomy, neurotomy, facet rhizotomy (C)</p> <p>IDET (I)</p> <p>Lumbar fusion unless concomitant instability or deformity proven (C)</p> <p>Artificial disc replacement (I)</p> |
| Spinal Fractures | <p>Bed rest for unstable spinal fractures (I)</p> <p>Vertebroplasty and</p> | <p>Gabapentin for chronic radicular pain syndromes (I)</p> <p>Mattress firmness</p> | <p>Sacroiliac joint fusion surgery and other sacroiliac joint surgical procedures (I)</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------|---|---|---|
| | Recommended | No Recommendation | Not Recommended |
| | <p>kyphoplasty for vertebral body compression fractures among those with chronic or severe pain (I)</p> <p>NSAIDs (I)</p> <p>Acetaminophen for patients with contraindications for NSAIDs (C)</p> <p>Gabapentin for perioperative pain management (A)</p> <p>Limited use (2 to 3 weeks) of opioids with longer periods for more severe fractures (C)</p> <p>Skeletal muscle relaxants as 2nd- or 3rd-line agents for more severe pain (I)</p> <p>Alteration of sleep posture (I)</p> <p>Gradual introduction of aerobic exercises during and to facilitate recovery (I)</p> <p>Strengthening exercises after aerobic exercises instituted and after healed (I)</p> <p>Stretching exercises to regain normal range of motion (I)</p> | <p>(I)</p> <p>Other optimal sleeping surfaces (e.g., bedding, water beds, hammocks) (I)</p> <p>Infrared therapy for home use (I)</p> <p>Ultrasound (I)</p> <p>Neuroreflexotherapy (I)</p> <p>Botulinum injections (I)</p> | <p>Percutaneous discectomy (nucleoplasty), laser discectomy, and disc coblation therapy (B)</p> <p>Adhesiolysis (I)</p> <p>Bed rest for stable spinal fractures (I)</p> <p>Vitamins (I)</p> <p>Willow bark (salix) (I)</p> <p>Spiroflor (I)</p> <p>Complementary or alternative treatments or dietary supplements, etc., other than those discussed in chapter (I)</p> <p>Reflexology (I)</p> <p>Lumbar supports (I)</p> <p>Magnets (I)</p> <p>Routine use of cryotherapies in health care provider offices or home use of high-tech device (I)</p> <p>Diathermy (I)</p> <p>Low-level laser therapy (I)</p> <p>Mechanical devices for administering massage (I)</p> <p>Interferential therapy (I)</p> <p>Taping and kinesiotaping (I)</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|---------------------|--|--|--|
| | Recommended | No Recommendation | Not Recommended |
| | Inclusion of FABT during course of rehabilitation (I) | | Myofascial release (I) High-voltage galvanic (I) Iontophoresis (I) Adjustments/manipulation (I) Acupuncture (I) Aggressive stretching (I) Strengthening of abdominal muscles (I) |
| Sacroiliitis | Sacroiliac joint corticosteroid injections for specific known cause of sacroiliitis (C) NSAIDs (I) Acetaminophen if contraindications for NSAIDs (I) Skeletal muscle relaxants as 2nd- or 3rd-line agents for more severe pain (I) Alteration of sleep posture (I) Aerobic exercises (I) Strengthening exercises after aerobic exercises instituted (I) Stretching exercises to regain normal | Gabapentin (I) Mattress firmness (I) Other optimal sleeping surfaces (e.g., bedding, water beds, hammocks) (I) Infrared therapy for home use (I) Ultrasound (I) Neuroreflexotherapy (I) Botulinum injections (I) | Sacroiliac joint fusion surgery and other sacroiliac joint surgical procedures (I) Bed rest for stable spinal fractures (I) Vitamins (I) Willow bark (salix) (I) Spiroflor (I) Complementary or alternative treatments or dietary supplements, etc., other than those discussed in chapter (I) Reflexology (I) Lumbar supports (I) Magnets (I) Routine use of cryotherapies in health care provider offices or home use of high-tech device (I) |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|--------------------------|---|---|---|
| | Recommended | No Recommendation | Not Recommended |
| | <p>range of motion (I)</p> <p>Inclusion of FABT during course of rehabilitation (I)</p> | | <p>Diathermy (I)</p> <p>Low-level laser therapy (I)</p> <p>Mechanical devices for administering massage (I)</p> <p>Interferential therapy (I)</p> <p>Taping and kinesiotaping (I)</p> <p>Myofascial release (I)</p> <p>High-voltage galvanic (I)</p> <p>Iontophoresis (I)</p> <p>Adjustments/manipulation (I)</p> <p>Acupuncture (I)</p> <p>Aggressive stretching (I)</p> <p>Strengthening of abdominal muscles (I)</p> |
| Spondylolisthesis | <p>Lumbar fusion for isthmic spondylolisthesis (C)</p> <p>Lumbar fusion for degenerative spondylolisthesis (C)</p> <p>NSAIDs (I)</p> <p>Acetaminophen if contraindications for NSAIDs (I)</p> <p>Skeletal muscle relaxants as 2nd- or</p> | <p>Gabapentin (I)</p> <p>Mattress firmness (I)</p> <p>Other optimal sleeping surfaces (e.g., bedding, water beds, hammocks) (I)</p> <p>Infrared therapy for home use (I)</p> <p>Ultrasound (I)</p> <p>Neuroreflexotherapy (I)</p> | <p>Bed rest (I)</p> <p>Vitamins (I)</p> <p>Willow bark (salix) (I)</p> <p>Spiroflor (I)</p> <p>Complementary or alternative treatments or dietary supplements, etc., other than those discussed in chapter (I)</p> <p>Reflexology (I)</p> <p>Lumbar supports (I)</p> |

| Low Back Disorder | Treatment with Evidence Rating/Recommendation Level | | |
|-------------------|---|--------------------------|---|
| | Recommended | No Recommendation | Not Recommended |
| | 3rd-line agents for more severe pain (I) Alteration of sleep posture (I) Aerobic exercises (I) Strengthening and stabilization exercises after aerobic exercises instituted (I) Inclusion of FABT during course of rehabilitation (I) | Botulinum injections (I) | Magnets (I) Routine use of cryotherapies in health care provider offices or home use of high-tech device (I) Diathermy (I) Low-level laser therapy (I) Mechanical devices for administration of massage (I) Interferential therapy (I) Taping and kinesiotaping (I) Myofascial release (I) High-voltage galvanic (I) Iontophoresis (I) Adjustments/manipulation (I) Acupuncture (I) Aggressive stretching (I) Strengthening of abdominal muscles (I) |

Definitions:

Strength of Evidence Ratings

A: Strong evidence-base: Two or more high-quality studies.¹

B: Moderate evidence-base: At least one high-quality study or multiple moderate-quality studies² relevant to the topic and the working population.

C: Limited evidence-base: At least one study of moderate quality.

I: Insufficient evidence: Evidence is insufficient or irreconcilable.

¹For therapy and prevention, randomized controlled trials (RCTs) with narrow confidence intervals and minimal heterogeneity. For diagnosis and screening, cross sectional studies using independent gold standards. For prognosis, etiology or harms, prospective cohort studies with minimal heterogeneity.

²For therapy and prevention, a well-conducted review of cohort studies. For prognosis, etiology or harms, a well-conducted review of retrospective cohort studies or untreated control arms of RCTs.

| Recommendation Category | Evidence Rating | Description of Category |
|---|------------------------|--|
| Strongly Recommended | A | The intervention is strongly recommended for appropriate* patients. The intervention improves important health and functional outcomes based on high quality evidence, and the Evidence-based Practice Panel (EBPP) concludes that benefits substantially outweigh harms and costs. |
| Moderately Recommended | B | The intervention is recommended for appropriate patients. The intervention improves important health and functional outcomes based on moderate quality evidence that benefits substantially outweigh harms and costs. |
| Recommended | C | The intervention is recommended for appropriate patients. There is limited evidence that the intervention may improve important health and functional benefits. |
| Insufficient - Recommended (Consensus-based) | I | The intervention is recommended for appropriate patients and has nominal costs and essentially no potential for harm.** The EBPP feels that the intervention constitutes best medical practice to acquire or provide information in order to best diagnose and treat a health condition and restore function in an expeditious manner. The EBPP believes based on the body of evidence, first principles, and/or collective experience that patients are best served by these practices, although the evidence is insufficient for an evidence-based recommendation. |
| Insufficient - No Recommendation (Consensus-based) | I | The evidence is insufficient to recommend for or against routinely providing the intervention. The EBPP makes no recommendation. Evidence that the intervention is effective is lacking, of poor quality, or conflicting and the balance of benefits, harms, and costs cannot be determined. |

| Recommendation Category | Evidence Rating | Description of Category |
|---|------------------------|---|
| Insufficient – NOT Recommended (Consensus-based) | I | The evidence is insufficient for an evidence-based recommendation. The intervention is not recommended for appropriate patients because of high costs/high potential for harm to the patient. |
| NOT Recommended | C | Recommendation against routinely providing the intervention. The EBPP found at least moderate evidence that harms and costs exceed benefits based on limited evidence. |
| Moderately NOT Recommended | B | Recommendation against routinely providing the intervention to eligible patients. The EBPP found at least moderate evidence that the intervention is ineffective, or that harms or costs outweigh benefits. |
| Strongly NOT Recommended | A | Strong recommendation against providing the intervention to eligible patients. The EBPP found high quality evidence that the intervention is ineffective, or that harms or costs outweigh benefits. |

* "Appropriate" means meeting screening or preventive method entry criteria without contraindications, or having the appropriate diagnosis, indication, time frame, prior conservative testing or treatment, and lack of contraindications for the specific test or treatment.

** For example, would include acetaminophen and self-administered cold or heat treatments. Excludes all interventional treatments, manual adjustment, and prescription medications. Aggregate and individual harms and costs are considered.

CLINICAL ALGORITHM(S)

The following clinical algorithms are provided in the original guideline document:

- Master low back algorithm
- Initial evaluation of acute and subacute low back and radicular pain
- Initial and follow-up management of acute and subacute low back and radicular pain
- Evaluation of subacute or slow-to-recover patients with low back pain unimproved or slow to improve (symptoms >4 weeks)
- Surgical considerations for patients with anatomic and physiologic evidence of nerve root compression and persistent low back symptoms
- Further management of subacute low back pain
- Further management of chronic low back pain

EVIDENCE SUPPORTING THE RECOMMENDATIONS

TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS

The type of supporting evidence is identified and graded for each recommendation (see "Major Recommendations").

Where there was not quality evidence, guidance represents a consensus of the Evidence-based Practice Spine Panel.

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

POTENTIAL BENEFITS

- Improved efficiency of the diagnostic process
- Effective treatment resulting in symptom alleviation and cure

POTENTIAL HARMS

- False-positive or false-negative diagnostic tests
- Risks and complications of surgical procedures and imaging studies (e.g., infection, radiation)
- Side effects of medications and injection therapies

CONTRAINDICATIONS

CONTRAINDICATIONS

Implanted metallic-ferrous device and significant claustrophobia are contraindications for magnetic resonance imaging (MRI)

QUALIFYING STATEMENTS

QUALIFYING STATEMENTS

- Original data from high- or moderate-quality randomized controlled clinical trials or cross-over trials were relied upon to develop all evidence-based treatment guidance. Many "systematic" reviews, low-quality randomized controlled studies, other studies, and other guidelines for treatments are referenced and reviewed in the Appendix of the original guideline document. Most of these "systematic" reviews or other guidelines are not high-quality reviews and guidelines usually due to one or more errors (e.g., lack of defined methodology, incomplete database searches, lack of exhaustive searches, lack of clearly defined article grading, lack of consideration of the importance of study design, selective use of the studies and inadequate or incorrect interpretation of the studies' results). These errors may render the conclusions invalid. Aside from Cochrane reviews which are nearly always high quality, these reviews, other studies, and other guidelines were not relied upon for purposes of the development of this document's guidance on treatments.
- The American College of Occupational and Environmental Medicine (ACOEM) provides this segment of guidelines for practitioners and notes that decisions to adopt particular courses of actions must be made by trained practitioners on the basis of the available resources and the particular circumstances

presented by the individual patient. Accordingly, the ACOEM disclaims responsibility for any injury or damage resulting from actions taken by practitioners after considering these guidelines.

IMPLEMENTATION OF THE GUIDELINE

DESCRIPTION OF IMPLEMENTATION STRATEGY

An implementation strategy was not provided.

IMPLEMENTATION TOOLS

Clinical Algorithm

For information about [availability](#), see the "Availability of Companion Documents" and "Patient Resources" fields below.

INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

IOM CARE NEED

Getting Better
Living with Illness

IOM DOMAIN

Effectiveness
Patient-centeredness

IDENTIFYING INFORMATION AND AVAILABILITY

BIBLIOGRAPHIC SOURCE(S)

Hegmann KT, ed. Low back disorders. In: Glass LS, editor(s). Occupational medicine practice guidelines: evaluation and management of common health problems and functional recovery in workers. 2nd ed. Elk Grove Village (IL): American College of Occupational and Environmental Medicine (ACOEM); 2007. p. 366. [1310 references]

ADAPTATION

Not applicable: The guideline was not adapted from another source.

DATE RELEASED

1997 (revised 2007)

GUIDELINE DEVELOPER(S)

American College of Occupational and Environmental Medicine - Medical Specialty Society

SOURCE(S) OF FUNDING

American College of Occupational and Environmental Medicine

GUIDELINE COMMITTEE

American College of Occupational and Environmental Medicine Practice Guidelines Committee

COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE

Editor-in-Chief: Kurt T. Hegmann, MD, MPH, FACOEM, FACP

Evidence-based Practice Spine Panel Chair: James Talmage, MD

Evidence-based Practice Spine Panel Members: Roger Belcourt, MD, MPH, FACOEM; Jill Galper, PT, Med; Elizabeth Genovese, MD, MBA, FACOEM; Michael Goertz, MD, MPH, FACOEM; James Lessenger, MD, FACOEM; Tom Mayer, MD; Scott Morris, MD, MPH; Kathryn Mueller, MD, MPH, FACOEM; Jack Richman, MD, FACOEM, CCFP, DOHS; Russell Travis, MD; Michael S. Weiss, MD, MPH, FACOEM, FAAPMR, FAANEM

Panel Consultants: Timothy K. Behrens, PhD, CHES; M. Jann Dewitt, PhD; Steven D. Feinberg, MD, MPH; Corey D. Fox, PhD, ABPP; Arun Garg, PhD, CPE; Paul Hooper, DC, MPH, MS; Christy Porucznik, PhD, MSPH; William Tellin, DC, DABCO; Matthew S. Thiese, MSPH, PhD-C

Methodology Committee Consultant: John P. Holland, MD, MPH, FACOEM

Managing Editors: Production: Marianne Dreger, MA; Research: Julie A. Ording, MPH; Editorial Assistant: Debra M. Paddack

FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST

Roger Belcourt, MD, MPH (Panel Member)

Assistant Vice President, Pacific Coast Zone, Concentra

National, Regional, Local Committee Affiliations—President, Nevada Health Professional Assistance Foundation; Diversion Committee and Foundation Board Member; Nevada Institutional Review Board; Board of Directors, Western Occupational and Environmental Medical Association (WOEMA); and Chair, 2006 Western Occupational Health Conference

Guidelines Related Professional Activities—None

Research Grants/Other Support—None

Financial/Non-Financial Conflict of Interest—None

Jill Galper, PT, MEd (Panel Member)

Vice President, Clinical Program Development, IMX Medical Management Services

National, Regional, Local Committee Affiliations—Executive Committee, Southeast District, Pennsylvania Physical Therapy Association

Guidelines Related Professional Activities—Affiliate Editor, ACOEM's APG Insights; American Physical Therapy Association (APTA)

Representative for Occupational Medicine Practice Guidelines, 2nd Edition, 2004

Research Grants/Other Support—None

Financial/Non-Financial Conflict of Interest—None

Elizabeth Genovese, MD, MBA (Panel Member)

Medical Director, IMX Medical Management Services; Adjunct Assistant Professor of Medicine, Associated Faculty, University of Pennsylvania School of Medicine

National, Regional, Local Committee Affiliations—Board of Directors, American Academy of Disability Evaluating Physicians; Advisory Committee, Athena Institute for Women's Wellness; Board of Directors, Philadelphia OEMS; Committee on Coding and Classification, ACOEM; Committee on Return to Work, ACOEM; Evidence Based Practice Committee, ACOEM; AMA CPT Advisory Committee, ACOEM Representative; Director, "Musculoskeletal Diagnosis and Treatment" course, ACOEM; Co-Director, "Clinical Guidelines" course, ACOEM

Guidelines Related Professional Activities—Member, Evidence Based Practice Committee, Occupational Medicine Practice Guidelines, 2nd Edition, 2004; Editor, ACOEM's APG Insights; Section Reviewer, AMA Guides to the Evaluation of Permanent Impairment, 6th Edition

Research Grants/Other Support—None

Financial/Non-Financial Conflict of Interest—None

Michael Goertz, MD, MPH (Panel Member)

Department Chair, Occupational Medicine, Park Nicollet Clinic

National, Regional, Local Committee Affiliations—Medical Services Review Board, DOL State of Minnesota

Guidelines Related Professional Activities—Guidelines Panel Member, Disability Prevention/Management, ACOEM; Member, Evidence Based Practice Committee, ACOEM Occupational Medicine Practice Guidelines, 2nd Edition, 2004

Research Grants/Other Support—None

Financial/Non-Financial Conflict of Interest—None

Kurt Hegmann, MD, MPH (Editor-in-Chief)

Associate Professor and Center Director, Rocky Mountain Center for Occupational and Environmental Health, University of Utah

National, Regional, Local Committee Affiliations—Member, Ergonomics Committee (Chair 2001-05), ACOEM; Board of Trustees, American Board of Preventive Medicine (Chair, Examination Committee); and Chair, Federal Motor Carrier Safety Administration's Medical Review Board

Guidelines Related Professional Activities—Chair, Evidence Based Practice Committee (update of 2nd Edition), ACOEM; Member, Council on Scientific Affairs (2001-05), ACOEM

Research Grants/Other Support—NIOSH (CDC) Training grants and research grants primarily on the epidemiology of musculoskeletal disorders (e.g., CTS, shoulder tendinosis, LBP) and truck driver safety; and a grant from the Utah Labor Commission studying cancers among firefighters and police officers

Financial/Non-Financial Conflict of Interest—Honoraria: Teaching honoraria from various courses, mostly ACOEM-related; Consultations: Consulting with companies regarding how to reduce work-related injuries, causation and apportionment of injuries and consultations with unions regarding return to work, work restrictions and work-relatedness on injuries; Clinical: Primary, secondary and tertiary clinical management of occupational injuries and diseases

John P. Holland, MD, MPH (Methodology Committee Consultant)

Principal, Holland Associates, Inc., Seattle, Washington; Clinical Assistant Professor, Department of Environmental and Occupational Health Sciences and Department of Orthopaedics, University of Washington, Seattle

National, Regional, Local Committee Affiliations—Board of Directors (1994-2005), ACOEM; President (2003-04), ACOEM; Chair, Council on External Affairs (2005-06), ACOEM

Guidelines Related Professional Activities—Chair, Practice Guidelines Committee (1998-2000), ACOEM; Chair, Practice Guidelines Steering Committee (2003-06), ACOEM; Methodology Advisory Group and Peer Reviewer (2001-04), Occupational Medicine Practice Guidelines, 2nd Edition, ACOEM; Head, Research Team (1991-94), Clinical Practice Guidelines on Low Back in Adults, AHCPR; Head, Research Team (1998-2002), Clinical Practice Guidelines for Young Children with Developmental Disabilities, New York State Department of Health

Research Grants/Other Support—None

Financial/Non-Financial Conflict of Interest—None

James Lessenger, MD (Panel Member)

Private Practice; Disability Examiner, Department of Social Services, State of California; Consultant, Toxcenter, Northridge Hospital Medical Center; Consultant, Medical Board of California; Lecturer, Occupational Medicine Residency, University of California, San Francisco

National, Regional, Local Committee Affiliations—Board Member, Benicia Historical Museum; Editorial Board, Journal of Agromedicine; Western Occupational Medicine Association

Guidelines Related Professional Activities—None

Research Grants/Other Support—None

Financial/Non-Financial Conflict of Interest—None

Tom Mayer, MD (Panel Member)

Private Practice: Medical Director, Productive Rehabilitation Institute of Dallas for Ergonomics (PRIDE), Dallas, TX; Clinical Professor of Orthopedic Surgery, University of Texas Southwestern Medical Center

National, Regional, Local Committee Affiliations—Editorial Board and Founding Editor, The Spine Journal; Board of Associate Editors, SPINE; Interdisciplinary Program Work Group, Division of Workers' Compensation/Texas Department of Insurance; International Editorial Board, Isokinetics and Exercise Science; Editorial Board, The Back Letter; Board of Trustees, Dallas Opera Association

Guidelines Related Professional Activities—Musculoskeletal Section Editor and Spine Chapter Author, Executive Editorial Board, 6th Edition, AMA Guides to the Evaluation of Permanent Impairment; Editorial Advisory Board, Official Disability Guidelines (ODG); AMA Guides Newsletter Advisory Board; Co-Chair, North American Spine Society, Spine Treatment Guideline (1996-04); Co-Chairman, Texas Spine Treatment Guideline Work Group (1990-95)

Research Grants/Other Support—None

Financial/Non-Financial Conflict of Interest—None

Scott Morris, MD, MPH (Panel Member)

Area Medical Director, Chicago, Concentra Medical Centers

National, Regional, Local Committee Affiliations—None

Guidelines Related Professional Activities—None

Research Grants/Other Support—None

Financial/Non-Financial Conflict of Interest—None

Kathryn Mueller, MD, MPH (Panel Member)

Medical Director, Colorado Division of Workers Compensation, Colorado Department of Labor; Professor of Surgery, Division of Emergency Medicine and Department of Preventive Medicine, University of Colorado Health Science Center; UCHSC Graduate School Faculty

National, Regional, Local Committee Affiliations—International Association of Injury and Accident Boards and Commissions Committee on Occupational Health and Disability Management; Bylaws Revision Committee, Committee on Workers' Compensation, Committee on Evidence Based Medicine, Steering Committee for Revision of ACOEM Guidelines, ACOEM; Task Force on Chronic Pain, Task Force Complex Regional Pain Syndrome/Reflex Sympathetic Dystrophy, Upper Extremity Task Force on Carpal Tunnel, Cumulative Trauma Disorder and Thoracic Outlet Syndrome, Task Force on Lumbar and Cervical Spine and Lower Extremity, Colorado Division of Workers' Compensation; Workers' Compensation Personal Injury and Workers' Compensation Committee, Colorado Medical Society; Preventive Medicine Residency Advisory Committee, Occupational Medicine Residency Advisory Committee, MSPH Policy Committee, MSPH Curriculum Committee, University of Colorado School of Medicine

Guidelines Related Professional Activities—Section Editor, AMA Guides to Evaluation of Permanent Impairment, 6th Edition; LBP Guideline Subcommittee, American Pain Society/American College of Physicians; Guidelines for State of Colorado; Editorial Board, AMA Guides Newsletter; Adviser/Reviewer, Medical Disability Advisor, 3rd Edition

Research Grants/Other Support—NIOSH Training Grant for Occupational Medicine Residencies, University of Colorado Health Sciences Department of Preventive Medicine

Financial/Non-Financial Conflict of Interest—Received less than \$2,000 in honorarium for teaching engagements for ACOEM

Jack Richman, MD (Panel Member)

Executive Vice President and Medical Director, AssessMed Inc.; and President, AssessMed Quality Review

National, Regional, Local Committee Affiliations—Chair, Research Committee of the Canadian Institute for the Relief of Pain and Disability (CIRPD)

Guidelines Related Professional Activities—Member, Guidelines Committee, ACOEM (2nd Edition); Ontario Government Occupational Disease Panel for the Workplace Safety and Insurance Board; and Chair, Standards Committee, Canadian Society of Medical Evaluators

Research Grants/Other Support—None

Financial/Non-Financial Conflict of Interest—None

James B. Talmage, MD (Chair of Panel)

Occupational Health Center LLC, Cookeville, Tennessee

National, Regional, Local Committee Affiliations—Member, Return to Work Committee, ACOEM; Examination Committee, ABIME; Editorial Advisory Board, Tennessee Workers' Comp Reporter; Editorial Board, Tennessee Medicine, Tennessee Medical Association

Guidelines Related Professional Activities—Chair, Spine Panel, Occupational Medicine Practice Guidelines; (update to 2nd Edition); Associate Editor, APG Insights, ACOEM; Associate Editor, AMA Guides Newsletter; Section Editor, Medical Disability Advisor, 3rd, 4th, 5th Editions; Reviewer, Guides to the Evaluation of Permanent Impairment, 5th Ed.

Research Grants/Other Support—None

Financial/Non-Financial Conflict of Interest—None

Russell Travis, MD (Panel Member)

Associate Medical Director, Cardinal Hill Rehabilitation Hospital; Voluntary Faculty, University of Kentucky Physical Medicine and Rehabilitation Department

National, Regional, Local Committee Affiliations—Board of Directors, American Academy of Disability Evaluating Physicians; Task Force on Health Care and Access and Affordability, Commonwealth of Kentucky; Board of Directors, Kentucky Foundation for Medical Care; Committee on Professional Liability Insurance, Committee on National Legislative Activities, Committee on Medical Insurance and Prepayment Plans, Kentucky Medical Association; Regional Translational Research Committee and Co-chair Selection Committee for Endowed Research Chair; Board of Directors, Kentucky Medical Insurance Company; Joint Washington Committee of Neurosurgery; Joint Section on Disorders of the Spine and Peripheral Nerves and Spine Task Force

Guidelines Related Professional Activities—Chapter Work Group, AMA's Guides to the Evaluation of Permanent Impairment, 6th Ed.; Editorial Advisory Board, Official Disability Guidelines, ODG Treatment; Advisory Board, ACOEM's APG Insights

Research Grants/Other Support—None

Financial/Non-Financial Conflict of Interest—None

Michael S. Weiss, MD, MPH (Panel Member)

Medical Director, Occupational Health Services, St. Luke's Regional Medical Center; Assistant Clinical Professor, Department of Rehabilitation Medicine/Department of Environmental and Occupational Medicine, University of

Washington School of Medicine; Medical Director, Idaho Power Company; Medical Director, Paradigm Health Corporation; Medical Consultant, Idaho State Insurance Fund

National, Regional, Local Committee Affiliations—Board, Northwest Occupational and Environmental Medical Association; Resident Advisory Committee, University of Washington, Department of Environmental and Occupational Medicine

Guidelines Related Professional Activities—None

Research Grants/Other Support—None

Financial/Non-Financial Conflict of Interest—None

GUIDELINE STATUS

This is the current release of the guideline.

This guideline updates a previous version: Low back complaints. In: Glass LS, editor(s). Occupational medicine practice guidelines: evaluation and management of common health problems and functional recovery in workers. 2nd ed. Elk Grove Village (IL): American College of Occupational and Environmental Medicine (ACOEM); 2004. p. 286-326.

The ACOEM *Guidelines* are currently being updated on a 3-year rolling process.

GUIDELINE AVAILABILITY

Print copies are available from ACOEM, 25 Northwest Point Boulevard, Suite 700, Elk Grove Village, IL 60007; Phone: 847-818-1800. To order a subscription to the online version, call 800-441-9674 or visit <http://www.acoempracguides.org/>.

AVAILABILITY OF COMPANION DOCUMENTS

None available

PATIENT RESOURCES

None available

NGC STATUS

This NGC summary was completed by ECRI on May 31, 2006. The information was verified by the guideline developer on November 3, 2006. This summary was updated by ECRI Institute on July 29, 2008. The updated information was verified by the guideline developer on August 7, 2008.

COPYRIGHT STATEMENT

The American College of Occupational and Environmental Medicine, the signator of this license, represent and warrant that they are the publisher of the guidelines and/or possess all rights necessary to grant the license rights to AHRQ and its agents.

DISCLAIMER

NGC DISCLAIMER

The National Guideline Clearinghouse™ (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at <http://www.guideline.gov/about/inclusion.aspx>.

NGC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI Institute, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.

© 1998-2008 National Guideline Clearinghouse

Date Modified: 10/6/2008

