



Complete Summary

GUIDELINE TITLE

Urinary incontinence.

BIBLIOGRAPHIC SOURCE(S)

Dowling-Castronovo A, Bradway C. Urinary incontinence. In: Mezey M, Fulmer T, Abraham I, Zwicker DA, editor(s). Geriatric nursing protocols for best practice. 2nd ed. New York (NY): Springer Publishing Company, Inc.; 2003. p. 83-98. [26 references]

GUIDELINE STATUS

Note: This guideline has been updated. The National Guideline Clearinghouse (NGC) is working to update this summary.

COMPLETE SUMMARY CONTENT

SCOPE
METHODOLOGY - including Rating Scheme and Cost Analysis
RECOMMENDATIONS
EVIDENCE SUPPORTING THE RECOMMENDATIONS
BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS
QUALIFYING STATEMENTS
IMPLEMENTATION OF THE GUIDELINE
INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT
CATEGORIES
IDENTIFYING INFORMATION AND AVAILABILITY
DISCLAIMER

SCOPE

DISEASE/CONDITION(S)

Urinary incontinence

GUIDELINE CATEGORY

Evaluation
Management
Treatment

CLINICAL SPECIALTY

Geriatrics
Nursing

INTENDED USERS

Advanced Practice Nurses
Nurses

GUIDELINE OBJECTIVE(S)

- To present a nursing standard of practice protocol for urinary incontinence (UI)
- To discuss the transient and established etiologies of UI
- To describe the core components of a nursing assessment for UI in hospitalized elders
- To identify major treatment strategies for UI
- To provide indications for indwelling catheter use

TARGET POPULATION

Older adults hospitalized for acute care

INTERVENTIONS AND PRACTICES CONSIDERED

Evaluation for Urinary Incontinence (UI)

1. Review medical and social history
2. Question patient/caregiver about possible urinary incontinence including time of onset, frequency and severity. Inquire about symptoms such as nocturia, hematuria, and hesitancy. Determine patient's current management strategy
3. Perform abdominal, genital, rectal, and skin examination
Valsalva maneuver for women (if not medically contraindicated)
4. Determine presence/absence of indwelling catheter. If present, determine whether use is indicated
5. Complete functional, environmental, and mental status assessment
6. Review medications and diagnostic test results
 - Urinalysis and urine cultures
 - Bladder sonography or catheter insertion to evaluate post-void residual urine
7. Complete bladder diary or voiding record
Staff training in use/interpretation of bladder diary

Treatment/Management

1. Scheduled voiding or habit training
2. Patient/caregiver, nursing staff education regarding urinary incontinence
 - Crede maneuver (deep suprapubic palpation)
 - Double voiding technique
 - Pelvic muscle exercises
3. Modification of environment to facilitate continence
 - Call bells within reach

- Commode, urinal, bedpan
- 4. Nutrition/hydration management
- 5. Referral to interdisciplinary team members (e.g., pharmacology, surgery, therapy) as indicated
- 6. Catheterization only as indicated
- 7. Avoidance of medications that may contribute to UI

MAJOR OUTCOMES CONSIDERED

- Number of episodes of urinary incontinence (UI)
- Number of complications associated with UI

METHODOLOGY

METHODS USED TO COLLECT/SELECT EVIDENCE

Searches of Electronic Databases

DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE

Medline, CINAHL, and MDConsult were the databases used.

NUMBER OF SOURCE DOCUMENTS

Not stated

METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE

Not stated

RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE

Not applicable

METHODS USED TO ANALYZE THE EVIDENCE

Review

DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE

Not stated

METHODS USED TO FORMULATE THE RECOMMENDATIONS

Not stated

RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS

Not applicable

COST ANALYSIS

A formal cost analysis was not performed and published cost analyses were not reviewed.

METHOD OF GUIDELINE VALIDATION

Internal Peer Review

DESCRIPTION OF METHOD OF GUIDELINE VALIDATION

Not stated

RECOMMENDATIONS

MAJOR RECOMMENDATIONS

Note: This guideline has been updated. The National Guideline Clearinghouse (NGC) is working to update this summary. The recommendations that follow are based on the previous version of the guideline.

Assessment Parameters

- Document the presence/absence of urinary incontinence (UI) for all patients on admission.
- Document the presence/absence of an indwelling urinary catheter. Determine appropriate indwelling catheter use: severely ill patients, patient with Stage III to IV pressure ulcers of the trunk, urinary retention unresolved by other interventions.
- For patients who are incontinent:
 - Determine whether the problem is transient, established, or both.
 - Identify and document the possible etiologies of the UI.
 - Elicit assistance with assessment and management from interdisciplinary team members.

Nursing Care Strategies

- General principals that apply to prevention and management of all forms of UI:
 - Identify and treat causes of transient UI.
 - Identify and continue successful pre-hospital management strategies for established UI.
 - Complete bladder diary.
 - Develop an individualized plan of care using data obtained from the history and physical examination, and in collaboration with other team members.
 - Avoid medications that may contribute to UI.
 - Avoid indwelling urinary catheters whenever possible.

- Monitor fluid intake and maintain an appropriate hydration schedule.
- Modify the environment to facilitate continence.
- Provide patients with usual undergarments in expectation of continence, if possible.
- Prevent skin breakdown by providing immediate cleansing after an incontinent episode and utilizing barrier ointments.
- Use absorbent products judiciously.
- Strategies for specific problems:

Stress UI

- Teach pelvic muscle exercises (PME).
- Provide toileting assistance and bladder training.
- Consider referral to other team members if pharmacologic or surgical therapies are warranted.

Urge UI

- Implement bladder training or habit training.
- Teach pelvic muscle exercises to be used in conjunction with the above strategy.
- Consider referral to other team members if pharmacologic therapy is warranted.
- Initiate referrals for those patients who do not respond to the above.

Overflow UI

- Allow sufficient time for voiding.
- Instruct patients in double voiding and Crede maneuver.
- Consider use of external collection devices for men.
- Provide sterile intermittent or indwelling catheterization.
- Initiate referrals to other team members for those patients requiring pharmacologic or surgical intervention.

Functional UI

- Provide scheduled voiding or habit training.
- Provide adequate fluid intake.
- Collaborate with other team members to eliminate any medications adversely affecting continence.
- Refer for physical and occupational therapy.
- Modify environment to be conducive to maintaining independence with continence.

Follow-up to Monitor the Condition

- Provide patient/caregiver discharge teaching regarding outpatient referral and management.
- Incorporate continuous quality improvement (CQI) criteria into existing program.

- Identify areas for improvement and enlist interdisciplinary assistance in devising strategies for improvement.

CLINICAL ALGORITHM(S)

None provided

EVIDENCE SUPPORTING THE RECOMMENDATIONS

TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS

The type of supporting evidence is not specifically stated for each recommendation.

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

POTENTIAL BENEFITS

Patients Will Demonstrate:

Fewer or no episodes of urinary incontinence (UI) or complications associated with UI

Health Care Providers Will Demonstrate:

- Documented continence status at admission and throughout hospital stay
- Interdisciplinary expertise and interventions to assess and manage UI during hospitalization
- Inclusion of UI in discharge planning needs and referral as indicated

Institution Will Demonstrate:

- Decreased incidence and prevalence of acute UI
- Hospital policies requiring assessment and documentation of continence status
- Access to the Agency for Health Care Policy and Research (AHCPR) Guidelines for Managing Acute and Chronic UI
- Administrative support and ongoing education regarding assessment and management of UI for staff

POTENTIAL HARMS

Urinary catheterization can be associated with urinary tract infections. Patients requiring indwelling urinary catheters may have a higher incidence rate of infection than patients requiring sterile intermittent catheterization.

QUALIFYING STATEMENTS

QUALIFYING STATEMENTS

Decisions regarding catheterization require careful consideration of the benefits and burdens associated with their use.

IMPLEMENTATION OF THE GUIDELINE

DESCRIPTION OF IMPLEMENTATION STRATEGY

An implementation strategy was not provided.

INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

IOM CARE NEED

Getting Better

IOM DOMAIN

Effectiveness
Patient-centeredness

IDENTIFYING INFORMATION AND AVAILABILITY

BIBLIOGRAPHIC SOURCE(S)

Dowling-Castronovo A, Bradway C. Urinary incontinence. In: Mezey M, Fulmer T, Abraham I, Zwicker DA, editor(s). Geriatric nursing protocols for best practice. 2nd ed. New York (NY): Springer Publishing Company, Inc.; 2003. p. 83-98. [26 references]

ADAPTATION

Not applicable: The guideline was not adapted from another source.

DATE RELEASED

2003

GUIDELINE DEVELOPER(S)

Hartford Institute for Geriatric Nursing - Academic Institution

GUIDELINE DEVELOPER COMMENT

The guidelines were developed by a group of nursing experts from across the country as part of the Nurses Improving Care for Health System Elders (NICHE) project, under sponsorship of The John A. Hartford Foundation Institute for Geriatric Nursing.

SOURCE(S) OF FUNDING

Supported by a grant from The John A. Hartford Foundation.

GUIDELINE COMMITTEE

Not stated

COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE

Primary Authors: Annemarie Dowling-Castronovo, APRN-BC, Gerontological Nurse Practitioner, New York University, The Steinhardt School of Education's Division of Nursing; Christine Bradway, PhD, APRN-BC, Gerontologic Nurse Practitioner, University of Pennsylvania School of Nursing

FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST

Not stated

GUIDELINE STATUS

Note: This guideline has been updated. The National Guideline Clearinghouse (NGC) is working to update this summary.

GUIDELINE AVAILABILITY

Electronic copies of the updated guideline: Available from the [Hartford Institute of Geriatric Nursing \(HIGN\) Web site](#).

AVAILABILITY OF COMPANION DOCUMENTS

None available

PATIENT RESOURCES

None available

NGC STATUS

This summary was completed by ECRI on February 2, 2004. The information was verified by the guideline developer on February 26, 2004.

COPYRIGHT STATEMENT

This NGC summary is based on the original guideline, which is subject to the guideline developer's copyright restrictions.

DISCLAIMER

NGC DISCLAIMER

The National Guideline Clearinghouse™ (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at <http://www.guideline.gov/about/inclusion.aspx>.

NGC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI Institute, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.

© 1998-2008 National Guideline Clearinghouse

Date Modified: 10/13/2008

