



## Complete Summary

---

### GUIDELINE TITLE

Surgical treatment of disease and injuries of the spleen.

### BIBLIOGRAPHIC SOURCE(S)

Society for Surgery of the Alimentary Tract (SSAT). Surgical treatment of disease and injuries of the spleen. Manchester (MA): Society for Surgery of the Alimentary Tract (SSAT); 2004 Feb 21. 3 p. [5 references]

### GUIDELINE STATUS

This is the current release of the guideline.

## COMPLETE SUMMARY CONTENT

SCOPE  
METHODOLOGY - including Rating Scheme and Cost Analysis  
RECOMMENDATIONS  
EVIDENCE SUPPORTING THE RECOMMENDATIONS  
BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS  
QUALIFYING STATEMENTS  
IMPLEMENTATION OF THE GUIDELINE  
INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT  
CATEGORIES  
IDENTIFYING INFORMATION AND AVAILABILITY  
DISCLAIMER

## SCOPE

### DISEASE/CONDITION(S)

Disease or injuries of the spleen:

- Traumatic injury to spleen
- Intraoperative splenic injury
- Hematologic diseases including immune thrombocytopenic purpura (ITP); hereditary spherocytosis; thalassemia major; certain forms of autoimmune hemolytic anemia unresponsive to medical management; myeloproliferative disorders; thrombotic thrombocytopenic purpura (TTP); and hairy-cell leukemia unresponsive to other treatment strategies
- Splenic abscesses, cysts, sinistral portal hypertension secondary to isolated splenic vein thrombosis or obstruction, splenic mass presumed to be a primary or undiagnosed neoplasm, malignancy in an adjacent organ

## **GUIDELINE CATEGORY**

Evaluation  
Treatment

## **CLINICAL SPECIALTY**

Family Practice  
Gastroenterology  
Hematology  
Internal Medicine  
Oncology  
Pediatrics  
Surgery

## **INTENDED USERS**

Physicians

## **GUIDELINE OBJECTIVE(S)**

To guide primary care physicians to the appropriate utilization of surgical procedures on the alimentary tract or related organs

## **TARGET POPULATION**

Patients (adults and children) with disease or injuries of the spleen

## **INTERVENTIONS AND PRACTICES CONSIDERED**

### **Evaluation**

1. Computed tomography (CT) scanning
2. Ultrasound

### **Treatment**

1. Non-operative support with in-hospital observation
2. Splenectomy (open or laparoscopic)
3. Spleen salvage
  - Suture plication to achieve hemostasis
  - Topical hemostatic agents (including absorbable mesh)
  - Electrocautery
  - Argon beam coagulation
4. Prophylaxis against post-splenectomy sepsis, including:
  - Daily penicillin (certain pediatric patients)
  - Pneumococcal vaccine using Pneumovax (non-elective splenectomy)
  - Vaccination against pneumococcus, *Haemophilus influenza*, meningococcus (before elective splenectomy)

## **MAJOR OUTCOMES CONSIDERED**

- Mortality rate after splenectomy
- Postoperative complications of splenectomy
- Late sequelae related to splenectomy, including post-splenectomy sepsis

## METHODOLOGY

### **METHODS USED TO COLLECT/SELECT EVIDENCE**

Searches of Electronic Databases

### **DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE**

Not stated

### **NUMBER OF SOURCE DOCUMENTS**

Not stated

### **METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE**

Not stated

### **RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE**

Not applicable

### **METHODS USED TO ANALYZE THE EVIDENCE**

Review

### **DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE**

Not applicable

### **METHODS USED TO FORMULATE THE RECOMMENDATIONS**

Expert Consensus

### **DESCRIPTION OF METHODS USED TO FORMULATE THE RECOMMENDATIONS**

The Society for Surgery of the Alimentary Tract (SSAT) guidelines are based on statements and recommendations that were overwhelmingly supported by clinical evidence. Each represents a consensus of opinion and is considered a reasonable plan for a specific clinical condition.

(See companion document: Gadacz TR, Traverso LW, Fried GM, Stabile B, Levine BA. Practice guidelines for patients with gastrointestinal surgical diseases. J Gastrointest Surg 1998;2:483-484.)

## **RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS**

Not applicable

## **COST ANALYSIS**

A formal cost analysis was not performed and published cost analyses were not reviewed.

## **METHOD OF GUIDELINE VALIDATION**

Internal Peer Review

## **DESCRIPTION OF METHOD OF GUIDELINE VALIDATION**

The guidelines were reviewed by several committee members and then by the entire committee on several occasions. Each guideline was then sent back to the original author for final comment and reviewed again by the committee. Each guideline was approved by the Board of Trustees of the Society for Surgery of the Alimentary Tract and final comments were reviewed by the committee.

(See companion document: Gadacz TR, Traverso LW, Fried GM, Stabile B, Levine BA. Practice guidelines for patients with gastrointestinal surgical diseases. J Gastrointest Surg 1998;2:483-484.)

# **RECOMMENDATIONS**

## **MAJOR RECOMMENDATIONS**

### **Indications for Splenectomy**

Traumatic injury to the spleen is no longer an immediate or mandatory indication for operation or splenectomy. Computed tomography (CT) scanning or ultrasound can accurately characterize splenic injury in patients with blunt trauma. Non-operative support with in-hospital observation for up to 5 days is indicated in children and adults with splenic injury and hemodynamic stability, provided there is no evidence of other intra-abdominal injuries that might require laparotomy. Accepted indications for operation in adults include hemodynamic instability, bleeding >1,000 mL, transfusion of more than 2 units of blood, or other evidence of ongoing blood loss. In children under 14 years old, more aggressive non-operative support is justified. When operative intervention is necessary, preservation of the spleen should be considered if bleeding can be controlled quickly and when there are no other life-threatening intra-abdominal injuries. Again, in children under 14 years of age, more aggressive attempts at intraoperative splenic salvage are justified. Splenic autotransplantation with a free-graft for maintenance of specific splenic immunity is of no proven value.

## **Iatrogenic (Intraoperative) Splenic Injury**

The spleen may be injured during the performance of intraperitoneal procedures, especially those involving the distal esophagus, stomach, distal pancreas, or splenic flexure of the colon. These injuries may occur directly from operative retractors or by traction on capsular adhesions leading to persistent bleeding. To avoid splenectomy, hemostasis should be attempted using suture plication, topical hemostatic agents (including absorbable mesh), electrocautery, or argon beam coagulation. However, if secure hemostasis is not possible before blood loss is sufficient to require blood transfusion, the patient is better managed by splenectomy than by repeated attempts at splenic salvage.

## **Hematologic Diseases**

Indications for splenectomy should be determined with the close cooperation of a hematologist/oncologist. Common indications include immune thrombocytopenic purpura (ITP), hereditary spherocytosis, thalassemia major, and certain forms of autoimmune hemolytic anemia unresponsive to medical management. Thrombotic thrombocytopenic purpura (TTP) and hairy-cell leukemia unresponsive to other treatment strategies are occasional indications for splenectomy.

Myeloproliferative disorders may lead to massive splenomegaly. Related symptoms may be best relieved by splenectomy although it does not usually alter overall survival. This information should be clearly discussed with the patient prior to operation, and they should be aware of the frequent requirement for blood or blood products when splenectomy is carried out for very large spleens. Massive splenomegaly may preclude a laparoscopic approach. In these circumstances an open or "hand-assisted" laparoscopic technique may be used. The operative morbidity and mortality rates are higher in these patients due to the hematologic comorbidity.

## **Other Indications for Splenectomy**

Less common indications for splenectomy include splenic abscesses, cysts, sinistral portal hypertension secondary to isolated splenic vein thrombosis or obstruction, or splenic mass presumed to be a neoplasm. Splenectomy is occasionally included in en bloc resection for malignancy in an adjacent organ. Distal pancreatectomy usually includes splenectomy if preservation of the splenic artery and vein is either contraindicated (malignancy) or technically impossible.

## **Post-Splenectomy Sepsis**

Most pediatricians believe that children who have undergone splenectomy before the age of 5 years should be treated with a daily dose of penicillin until the age of 10 years. The benefit of prophylactic penicillin is less clear in children over 5 years old and in adults. All patients who have undergone non-elective splenectomy should be immunized with Pneumovax (a non-viable pneumococcal vaccine). When planning elective splenectomy, patients should be immunized with Pneumovax, and against *Haemophilus influenza* and meningococcus, preferably two or more weeks before operation.

## **Qualifications for Performing Operations on the Spleen**

The qualifications of a surgeon performing any operative procedure should be based on training (education), experience, and outcomes. At a minimum, surgeons who are certified or eligible for certification by the American Board of Surgery, the Royal College of Physicians and Surgeons of Canada, or their equivalent should perform emergency and elective operations on the spleen. For laparoscopic splenic procedures, surgeons should have advanced laparoscopic training and expertise.

### **CLINICAL ALGORITHM(S)**

None provided

## **EVIDENCE SUPPORTING THE RECOMMENDATIONS**

### **TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS**

The type of supporting evidence is not specifically stated for each recommendation.

## **BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS**

### **POTENTIAL BENEFITS**

- Appropriate use of surgical treatment for diseases or injuries of the spleen
- Laparoscopic splenectomy, which is becoming increasingly common, appears to be safe and associated with less pain, shorter hospital stay, and more rapid convalescence than traditional open splenectomy.

### **POTENTIAL HARMS**

- Operative mortality for elective splenectomy is less than 1% except in patients with myeloproliferative disorders, who are at increased risk for postoperative hemorrhage. In trauma patients, the mortality rate for splenectomy depends upon the extent of other injuries. Postoperative complications of open splenectomy include pneumonia, thrombotic complications, wound infection, hernia formation, hemorrhage, subphrenic abscess, pancreatic abscess/fistula, pancreatic pseudocyst, and, rarely, gastric fistula/perforation. These potential complications also exist when using the laparoscopic approach, although wound complications consist primarily of herniation at trocar sites.
- Late sequelae related to splenectomy are much more common in children, especially those under 6 years old. Overwhelming post-splenectomy sepsis is a rare (less than 1%) but potentially fatal complication of splenectomy. It is much more common in children younger than age 6 who have not yet developed extra-splenic specific immunity to encapsulated organisms such as pneumococcus and meningococcus. Adults are susceptible to similar infections following splenectomy, but the incidence is likely much lower than in children.

## QUALIFYING STATEMENTS

### QUALIFYING STATEMENTS

These patient care guidelines were written for the primary care physicians on a variety of digestive diseases to assist on when to refer the patient for surgical consultation. Their goal is to guide primary care physicians to the appropriate utilization of surgical procedures on the alimentary tract or related organs, and they are based on critical review of the literature and expert opinion. Both of the latter sources of information result in a consensus that is recorded in the form of these Guidelines. The consensus addresses the range of acceptable clinical practice and should not be construed as a standard of care. These Guidelines require periodic revision to ensure that clinicians utilize procedures appropriately but the reader must realize that clinical judgment may justify a course of action outside of the recommendations contained herein.

## IMPLEMENTATION OF THE GUIDELINE

### DESCRIPTION OF IMPLEMENTATION STRATEGY

An implementation strategy was not provided.

### IMPLEMENTATION TOOLS

Foreign Language Translations

For information about [availability](#), see the "Availability of Companion Documents" and "Patient Resources" fields below.

## INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

### IOM CARE NEED

Getting Better  
Living with Illness

### IOM DOMAIN

Effectiveness

## IDENTIFYING INFORMATION AND AVAILABILITY

### BIBLIOGRAPHIC SOURCE(S)

Society for Surgery of the Alimentary Tract (SSAT). Surgical treatment of disease and injuries of the spleen. Manchester (MA): Society for Surgery of the Alimentary Tract (SSAT); 2004 Feb 21. 3 p. [5 references]

**ADAPTATION**

Not applicable: The guideline was not adapted from another source.

**DATE RELEASED**

2004 Feb 21

**GUIDELINE DEVELOPER(S)**

Society for Surgery of the Alimentary Tract, Inc - Medical Specialty Society

**SOURCE(S) OF FUNDING**

Society for Surgery of the Alimentary Tract, Inc

**GUIDELINE COMMITTEE**

Patient Care Committee

**COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE**

Not stated

**FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST**

Not stated

**GUIDELINE STATUS**

This is the current release of the guideline.

**GUIDELINE AVAILABILITY**

Electronic copies: Available from the [Society for Surgery of the Alimentary Tract, Inc. Web site](#).

Print copies: Available from the Society for Surgery of the Alimentary Tract, Inc., 900 Cummings Center, Suite 221-U, Beverly, MA 01915; Telephone: (978) 927-8330; Fax: (978) 524-0461.

**AVAILABILITY OF COMPANION DOCUMENTS**

The following is available:

- Society for Surgery of the Alimentary Tract (SSAT). Surgical treatment of disease and injuries of the spleen (Spanish version). Manchester (MA): Society for Surgery of the Alimentary Tract (SSAT); 2004 Feb 21. Electronic copies available from the [Society for Surgery of the Alimentary Tract, Inc. Web site](#).

- Gadacz TR, Traverso LW, Fried GM, Stabile B, Levine BA. Practice guidelines for patients with gastrointestinal surgical diseases. J Gastrointest Surg 1998;2:483-484.

Print copies: Available from the Society for Surgery of the Alimentary Tract, Inc., 900 Cummings Center, Suite 221-0, Beverly, MA 01915; Telephone: (978) 927-8330; Fax: (978) 524-8890.

## **PATIENT RESOURCES**

None available

## **NGC STATUS**

This NGC summary was completed by ECRI on October 19, 2004.

## **COPYRIGHT STATEMENT**

For terms governing downloading, use, and reproduction of these guidelines, please contact: [ssat@prri.com](mailto:ssat@prri.com).

## **DISCLAIMER**

### **NGC DISCLAIMER**

The National Guideline Clearinghouse™ (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at <http://www.guideline.gov/about/inclusion.aspx>.

NGC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI Institute, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.

© 1998-2008 National Guideline Clearinghouse

Date Modified: 11/3/2008

